

River export of bioavailable nutrients and labile organic carbon effects on estuarine nutrient stoichiometry and bacterioplankton nutrient limitation

Soares, Ana; Marchlewska, Joanna; Islam, Sohidul; Berggren, Martin

2014

## Link to publication

Citation for published version (APA): Soares, A., Marchlewska, J., Islam, S., & Berggren, M. (2014). River export of bioavailable nutrients and labile organic carbon: effects on estuarine nutrient stoichiometry and bacterioplankton nutrient limitation. Poster session presented at ASLO 2014 Ocean Sciences Meeting.

Total number of authors:

## General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study

- · You may not further distribute the material or use it for any profit-making activity or commercial gain You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

## Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Alves Soares AR, J Marchlewska, S Islam and M Berggren (poster). 2014. River export of bioavailable nutrients and labile organic carbon: effects on estuarine nutrient stoichiometry and bacterioplankton nutrient limitation. *AGU/ASLO 2014 Ocean Sciences Meeting*, presented on February 23-28, 2014, in Honolulu, Hawaii, USA

## **ABSTRACT**

Nutrient control of planktonic processes is a fundamental aspect of aquatic ecology that is not well understood in northern ecosystems. Boreal forests and wetlands export large amounts of dissolved organic carbon (DOC), total nitrogen (TN) and phosphorus (TP) via river runoff, but a highly variable portion of the nutrient pools is bioavailable. We used a bioassay method to simultaneously measure the bioavailability of TN, TP and DOC in upland headwater sources, and in river mouths that represent the points of entry to the Baltic Sea. The stoichiometry of bioavailable elements was vastly different from the stoichiometry of bulk carbon and nutrient fractions. We show that TN, although being dominated by organic N, had a much higher degree of bioavailability than DOC and TP. Our preliminary results demonstrate that the stoichiometry of bioavailable elements can be used to accurately predict nutrient limitation of estuarine bacterioplankton, in situations where predictions based on bulk nutrients would fail. Our approach opens up new possibilities to assess the ecological status of fresh and coastal waters, and to evaluate the effects of nutrient loads on plankton communities.