



LUND UNIVERSITY

Radiations from a Water Jet Plasma Source

Fullagar, Wilfred; Uhlig, Jens; Walczak, Monika; Canton, Sophie; Wahlström, Claes-Göran; Sundström, Villy

2007

Document Version:
Other version

[Link to publication](#)

Citation for published version (APA):

Fullagar, W., Uhlig, J., Walczak, M., Canton, S., Wahlström, C.-G., & Sundström, V. (2007). *Radiations from a Water Jet Plasma Source*. 7.

Total number of authors:
6

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>


Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Wilfred Fullagar, Jens Uhlig, Monika Walczak, Sophie Canton, Claes-Göran Wahlström, Villy Sundström



X-ray shadow image of a leaf


Division of Chemical Physics
Division of Atomic Physics




X-ray shadow image of a fly head

Laser system

low power branch of LLC terawatt facility (June 2007)




Output beam:
800 nm, ≥ 35 fs, ≤ 300 mJ/pulse, 10 Hz, ~ 4 cm ϕ




Typical temporal contrast (courtesy Filip Lindau)

X-rays




Goals:

- In-house subpicosecond chemical structure dynamics, via
 - pump-probe EXAFS
 - pump-probe Laue crystallography






Shadow of ightbulb filament & 250µm Cu grid
Blurring \Rightarrow source ~ 30 µm



A typical setup, shield chamber removed

Source ambitions:	This source:
* Everyday local development access	Yes (LLC collaboration)
* Simplicity \rightarrow in-house development & maintenance	Yes
* Broadband radiation	Yes (Maxwellian, $T = 10\text{-}50 \times 10^6$ K)
* Sub-picosecond burst	Expected (believed limited by (jet ϕ)/(e- velocity), or laser pulse)
* No emission lines (\rightarrow detection & mechanism issues)	Yes (oxygen $K\alpha$ too soft for typical filter transmission)
* Adequate X-ray flux	Yes (EXAFS) but requires detection development
* jitter-free laser synchronisation	Yes
* Collimation	No (needed for Laue xtlgrphy; see electron beams, below)
* Small source size	Yes (multi-shot images $\Rightarrow \sim 30$ µm in target plane)




Adaptable to typical kHz laser


Handy bunch of numbers (corresp. 10 Hz, 1 atm He operation)

Energy/pulse (mJ)	optical pulse duration	N_0 (ph/[eV.sr.sec])	T (K)
40	~ 200 fs (optimum)	71500	17.4×10^6
1.6	~ 200 fs (optimum)	14702	12.7×10^6
40	~ 40 fs (shortest)	29200	11.5×10^6

Selected relevant literature:
* R.J. Tompkins et al. Rev.Sci. Inst 69(9), 3113 (1998)
* K. Eidmann et al. Europhys Lett., 55(3) 334 (2001)




Compressor scan: X-ray yield & autocorrelator FWHM overlaid



Single shot Ti $K\alpha$ XAS edge spectrum (5 µm Ti foil)

Add s-pol prepulse, several 10s ps early, 1:1 pwr ratio
 \Rightarrow X-ray-free, controllable preplasma (vast parameter space)


Electrons




Electron beams away from and through the water jet

Also! : dramatic change of interaction geometry for continuum X-rays
(typical literature $\Rightarrow 30^\circ - 60^\circ$ incidence for assorted K_α)


Clear flashes on GdOS screen after ~ 1 cm air
 \Rightarrow observe strong scatter by 50 µm Be window




Collimated:
including strong filter scatter & assoc 2ndary radiation $< 10^\circ$ FWHM




Generates 5.9keV $K\alpha$ emission line through 5 mm Ti X-ray attenuation foil.




Deflects like electron in magnetic field (detector view from source)



Prepulse delay scan through $t=0$. $I \rightarrow r \approx 2$ ns.
 \Rightarrow e-beam requires p-polarised pulse in preplasma.




Electron ranges (CSDA, ESTAR, NIST). ρ^{-1} (cm²/g):
 $H_2O = 1$; He = 5590, Be = 0.541, air = 769, Si = 0.429
Suggests observed beams are 20 – 200 keV.



Strong e- generation when s- & p- pulses briefest. Suggests:
* proton motion during p-pulse unimportant
* e- beam emission = non-resonant phenomenon
* e- beam pulse duration \leftarrow energy range, distance from src

Selected literature precedents:
* S. Bastiani et al. Phys. Rev. E, 56(6), 7179 (1997)
* X.-Y. Peng et al. Chin. Phys. Lett., 21(4), 693 (2004)


Possible future?



Selected relevant literature:
* A. Rousse et al. Phys. Rev. Lett., 93(13), 135005 (2004)
* W.P. Leemans et al. Nature Physics, October(2), 696 (2006)


Protons??

H^+ emission conceivable with long pulses, high NA focus
For example:




H^+ jets moving to left, PIC simulation
F. Brunel, Phys. Fluids, 31(9) 2716 (1988)

Simulated electron spectra with & without mobile H^+
P. Gibbon, PRL 73(5), 664 (1994)



Circuit observes all ionization in chamber \rightarrow strong signal when hitting jet.
But! : only partial correlation with X-ray yield... why?

Protons stopped in chamber by He (\rightarrow ionisation) : see below
However : 524 eV oxygen $K\alpha$ (1/e in He $\rightarrow 12$ cm), & possible other contributions



Proton range in He (p-STAR, NIST)
 $\rho^{-1} = 5990$ cm²/g
 $\Rightarrow H^+ < \sim 1$ MeV stopped in chamber