



LUND UNIVERSITY

A variant in the transcription factor 7-like 2 (TCF7L2) gene is associated with an increased risk of gestational diabetes mellitus.

Shaat, Nael; Lernmark, Åke; Ekholm, Ella; Ivarsson, Sten; Parikh, Hemang; Berntorp, Kerstin; Groop, Leif

Published in:
Diabetologia

DOI:
[10.1007/s00125-007-0623-2](https://doi.org/10.1007/s00125-007-0623-2)

2007

[Link to publication](#)

Citation for published version (APA):

Shaat, N., Lernmark, Å., Ekholm, E., Ivarsson, S., Parikh, H., Berntorp, K., & Groop, L. (2007). A variant in the transcription factor 7-like 2 (TCF7L2) gene is associated with an increased risk of gestational diabetes mellitus. *Diabetologia*, 50(5), 972-979. <https://doi.org/10.1007/s00125-007-0623-2>

Total number of authors:
7

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



LUND UNIVERSITY
Faculty of Medicine

LU:*research*

Institutional Repository of Lund University

This is an author produced version of a paper published in Diabetologia. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Citation for the published paper:

Shaath, N and Lernmark, A and Karlsson, E and Ivarsson, S and Parikh, H and Berntorp, K and Groop, L.

"A variant in the transcription factor 7-like 2 (TCF7L2) gene is associated with an increased risk of gestational diabetes mellitus"

Diabetologia, 2007, Vol: 50, Issue: 5, pp. 972-9.

<http://dx.doi.org/10.1007/s00125-007-0623-2>

Access to the published version may
require journal subscription.

Published with permission from: Springer Verlag

**A variant in the transcription factor 7-like 2 (TCF7L2) gene
is associated with an increased risk of gestational diabetes
mellitus**

N. Shaaf¹, Å. Lernmark^{1, 2}, E. Karlsson¹, S. Ivarsson³, H. Parikh¹, K. Berntorp¹, L. Groop^{1, 4}.

¹Department of Clinical Sciences/Diabetes & Endocrinology, Malmö University Hospital, Lund University, Malmö, Sweden.

²Robert H. Williams Laboratory, University of Washington, Seattle, WA, USA.

³Department of Paediatrics, Malmö University Hospital, Lund University, Malmö, Sweden.

⁴Department of Medicine, Helsinki University Hospital, Helsinki, Finland.

Corresponding author

N. Shaaf

Department of Clinical Sciences/Diabetes & Endocrinology

Malmö University Hospital

Lund University

Malmö, Sweden

E-mail: nael.shaat@med.lu.se

Received: 24 October 2006/ Accepted: 21 January 2007

Abstract

Aims/hypothesis Genetic and epidemiological studies suggest an association between gestational diabetes mellitus and type 2 diabetes. Both are polygenic multifactorial disorders characterised by beta cell dysfunction and insulin resistance. Our aim was to investigate whether common genetic variants that have previously been associated with type 2 diabetes or related phenotypes would also confer risk for gestational diabetes mellitus.

Methods In 1881 unrelated pregnant Scandinavian women (649 women with gestational diabetes mellitus, 1232 non-diabetic control subjects) we genotyped the transcription factor 7-like 2 (*TCF7L2* rs7903146), adiponectin (*ADIPOQ* +276G>T), peroxisome-proliferator activated receptor, gamma 2 (*PPARG* Pro12Ala), PPARG-coactivator, 1 alpha (*PPARGC1A* Gly482Ser), forkhead box C2 (*FOXC2* -512C>T) and β 3-adrenergic receptor (*ADRB3* Trp64Arg) polymorphisms using TaqMan allelic discrimination assay or RFLP.

Results The CC, CT and TT genotype frequencies of the *TCF7L2* rs7903146 variant differed significantly between women with gestational diabetes mellitus and control women (46.3, 43.6 and 10.1% vs 58.5, 35.3 and 6.2%, $p=3.7 \times 10^{-6}$, corrected p value [Pc] for multiple testing $P_c=2.2 \times 10^{-5}$). The T-allele was associated with an increased risk of gestational diabetes mellitus (odds ratio 1.49 [95% CI 1.28–1.75], $p=4.9 \times 10^{-7}$ [Pc=2.8x10⁻⁶]). Compared with wild-type CC-genotype carriers, heterozygous (CT-genotype) and homozygous (TT-genotype) carriers had a 1.6-fold (95% CI 1.26–1.93, $p=3.7 \times 10^{-5}$ [Pc=0.0002]) and a 2.1-fold (95% CI 1.41–2.99, $p=0.0001$ [Pc=0.0008]) increased risk of gestational diabetes mellitus, respectively. The other polymorphisms studied were not significantly associated with gestational diabetes mellitus (*ADIPOQ* +276G>T: 1.17 [1.01–1.36], $p=0.039$ [Pc=0.23]; *PPARG* Pro12Ala: 1.06 [0.87–1.29], $p=0.53$; *PPARGC1A* Gly482Ser: 0.96 [0.83–1.10], $p=0.54$; *FOXC2* -512C>T: 1.01 [0.87–1.16], $p=0.94$; and *ADRB3* Trp64Arg: 1.22 [0.95–1.56], $p=0.12$).

Conclusions/interpretation The *TCF7L2* rs7903146 variant is associated with an increased risk of gestational diabetes mellitus in Scandinavian women.

Keywords

Adiponectin, *ADRB3*, Association, *FOXC2*, GDM, Gestational diabetes mellitus, Polymorphism, *PPARG*, *PPARGC1A*, *TCF7L2*.

Abbreviations

ADIPOQ	adiponectin,
ADRB3	β 3-adrenergic receptor
DBS	dried blood spots
FOXC2	forkhead box C2
GCK	glucokinase
HNF1A	hepatocyte nuclear factor 1- α
KCNJ11	potassium inwardly rectifying channel subfamily J, member 11
MAF	minor allele frequency
OR	odds ratio
<i>P_c</i>	corrected p value
PPAR	peroxisome-proliferator activated receptor
PPARG	peroxisome-proliferator activated receptor, gamma 2
PPARGC1A	PPARG-coactivator 1 alpha
SNP	single nucleotide polymorphism
TCF1	transcription factor 1, hepatic; LF-B1, hepatic nuclear factor (HNF1), albumin proximal factor
TCF7L2	transcription factor 7-like 2

Introduction

Genetic and epidemiological studies suggest an association between gestational diabetes mellitus and type 2 diabetes [1, 2]. The prevalence of gestational diabetes mellitus is increasing in parallel with the increased prevalence of type 2 diabetes [1, 2]. The former is a common metabolic disorder of pregnancy, affecting at least 2% of Scandinavian women and 5 to 10% of Asian, Hispanic/Mexican–American and Arabian women [2]. Gestational diabetes mellitus occurs when pancreatic beta cells are unable to compensate for increased insulin resistance during pregnancy [3]. Both insulin deficiency and insulin resistance are considered to be heritable with heritability estimates for insulin secretion of 0.75 to 0.84 and for insulin sensitivity of 0.53 to 0.55 [4].

Studies have consistently shown that women with a family history of diabetes have an increased risk of gestational diabetes mellitus [5, 6]. In addition, we and others have demonstrated that gestational diabetes mellitus shares some genetic risk factors with type 2 diabetes [2]. For example, potassium inwardly rectifying channel subfamily J, member 11 (*KCNJ11* E23K), glucokinase (*GCK* -30G>A) and transcription factor 1, hepatic; LF-B1, hepatic nuclear factor (HNF1), albumin proximal factor (TCF1, also known as hepatocyte nuclear factor 1- α [*HNF1A* I27L]) polymorphisms, all of which have been associated with type 2 diabetes [7–9], also increase the risk of gestational diabetes mellitus with a modest effect size [10, 11].

In the present study, therefore, we continued to study common genetic variants that have previously been associated with type 2 diabetes or related phenotypes such as the metabolic syndrome. In order to provide sufficient power to detect a modest odds ratio (OR \geq 1.3), we selected five variants with a minor allele frequency (MAF) of at least 15%. These polymorphisms were the transcription factor 7-like 2 (*TCF7L2* rs7903146)

[12–19], adiponectin (*ADIPOQ* +276G>T) [20, 21], peroxisome-proliferator activated receptor (PPAR), gamma 2 (*PPARG* Pro12Ala) [22], PPARG-coactivator 1, alpha (*PPARGC1A* Gly482Ser) [23] and forkhead box C2 (*FOXC2* –512C>T) [24]. Despite its MAF ~10%, we also included the β 3-adrenergic receptor (*ADRB3* Trp64Arg) polymorphism, as its putative association with gestational diabetes mellitus has been addressed in a number of small studies with inconsistent results [25–28]. These polymorphisms were genotyped in a case–control study of 1881 unrelated pregnant Scandinavian women (649 with gestational diabetes mellitus, 1232 non-diabetic control subjects).

Subjects and methods

Study subjects

Since 1995, all pregnant women in southern Sweden (Skåne) have been routinely offered a 75-g OGTT at 27 to 28 weeks of pregnancy, irrespective of family history of diabetes or any other risk factor for gestational diabetes mellitus. Women with previous gestational diabetes mellitus or a family history of diabetes are offered an additional 75-g OGTT at 12 to 13 weeks. The tests are performed in the local Maternity Health Care clinics, using HemoCue devices (HemoCue, Ängelholm, Sweden) for capillary whole-blood analysis. Gestational diabetes mellitus is defined, according to the proposal by the Diabetic Pregnancy Study Group of the European Association for the Study of Diabetes [29], as a 2-hour capillary glucose concentration (double test) of at least 9 mmol/l. The compliance rate for OGTTs is approximately 90%.

We studied 1881 unrelated pregnant Scandinavian women (649 with gestational diabetes mellitus, 1232 pregnant non-diabetic control subjects). The diabetic women were recruited from Malmö or Lund University Hospitals between March 1996 and December 2003 (n=226) as well as from the women participating in the Diabetes Prediction in Skåne study, which is a prospective, longitudinal study for the prediction of type 1 diabetes in all newborns in southern Sweden during the period from September 2000 until August 2004 (n=423) [30]. All pregnant non-diabetic control subjects (n=1232) were drawn from the same study [30]. Both study groups, i.e women with gestational diabetes mellitus and control women, are considered to be genetically homogeneous since they have the same ethnic background (i.e. Scandinavian) and were recruited from the same place and during the same time period. The characteristics of the majority of participants in the present study have been reported

earlier [10]. Detailed phenotypic characteristics, including OGTT data, were available only for a small proportion of the women with gestational diabetes mellitus [31]. Informed voluntary consent was obtained from all study subjects. The study was approved by the ethics committee of Lund University.

Genetic analyses

Samples' collection Total DNA was isolated from peripheral blood lymphocytes from 226 women with gestational diabetes mellitus. In the other subjects, blood samples were collected as dried blood spots (DBS) on filters (Grade 2992 filters; Schleicher and Schuell, Dassel, Germany).

Genotyping using DNA Genotyping of *TCF7L2* (rs7903146), *ADIPOQ* +276G>T (rs1501299), *PPARGC1A* Gly482Ser (rs8192678), *FOXC2* -512C>T and *ADRB3* Trp64Arg (rs4994) was carried out using TaqMan allelic discrimination assay, whereas *PPARG* Pro12Ala (rs1801282) polymorphism was genotyped by TaqMan assay or RFLP as previously described [31]. TaqMan assay was carried out on an ABI Prism 7900 Sequence Detection System (Applied Biosystems, Foster City, CA, USA) using 2 µl of DNA (5–10 ng) according to the manufacturer's instructions. Primers and probes were designed using Assays-by-Design (Applied Biosystems), except for *FOXC2* -512C>T, which was ordered from MWG Biotech Scandinavia (Risskov, Denmark). The primers and probes used are listed in Electronic supplementary material (ESM) Table 1.

Genotyping using DBS Using the primers listed in ESM Table 2, a template PCR was initially carried out to amplify the region of interest where the single nucleotide polymorphism (SNP) is located. The template PCR was followed by TaqMan allelic

discrimination assay or RFLP. The primers used for the template PCRs were designed to be located outside the TaqMan primers and probes.

The template PCR was performed with an initial two cycles (4°C for 30 s, followed by 98°C for 3 min), followed by holding at 80°C while the PCR mix was added. Then the PCR was continued with initial denaturation (94°C for 5 min, except for *ADIPOQ* +276G>T at 96°C), followed by 45 cycles of denaturation (94°C for 30 s, except for *ADIPOQ* +276G>T at 96°C), annealing (30 s) and extension (72°C for 30–60 s), followed by final extension (72°C for 10 min). The reagents used for PCR amplification are listed in ESM Table 3.

The template PCR was followed by SNP genotyping, which was carried out using TaqMan allelic discrimination assay with 2 µl of the template PCR product according to the manufacturer's instructions. However, for some of the DBS samples, the *ADIPOQ* +276G>T polymorphism was genotyped by RFLP where 20 µl of the template PCR product was digested with Mva 1269I (Fermentas, St Leon-Rot, Germany) at 37°C for 4 h. PCR products were separated on 3% agarose gel (SeaKem, Rockland, ME, USA) and stained with ethidium bromide to visualise the fragments using ultra-violet light. The *PPARG* Pro12Ala polymorphism was also genotyped by RFLP for some of the DBS samples as previously described [31].

Genotyping quality control Genotyping success rate was similar for women with gestational diabetes and for control subjects. In the former the success rates were: 90.1% (*TCF7L2* rs7903146); 98.8% (*ADIPOQ* +276G>T); 98.1% (*PPARG* Pro12Ala); 99.2% (*PPARGC1A* Gly482Ser); 97.8% (*FOXC2* –512C>T); and 98.5% (*ADRB3* Trp64Arg). For control subjects the success rates were: 90.2% (*TCF7L2* rs7903146); 99.4% (*ADIPOQ* +276G>T); 100% (*PPARG* Pro12Ala); 99.4% (*PPARGC1A*

Gly482Ser); 97.7% (*FOXC2* -512C>T); and 99.6% (*ADRB3* Trp64Arg). Genotyping error rate was determined to be 0.4% using 1751 (15.5%) duplicate genotypes as well as 89 double samples (i.e. women with gestational diabetes mellitus who had both peripheral blood DNA and DBS or two DBS taken at different deliveries). All polymorphisms conformed to Hardy–Weinberg equilibrium (Chi-squared $p > 0.05$) in gestational diabetes mellitus and control groups. These quality control measures indicate that genotyping results of all SNPs are reliable for analyses.

Statistical analyses

ANOVA was used to test the significance of difference in continuous variables such as age between gestational diabetes mellitus and control groups. Age was presented as mean \pm SEM. Chi-squared analysis was used to test for difference in genotype and allele frequencies between gestational diabetes mellitus and control groups. Logistic regression analysis was used to calculate the age-adjusted and/or crude ORs and 95% CIs for the polymorphisms. Statistical analyses were performed using the Number Cruncher Statistical Systems (NCSS, Kaysville, UT, USA). Bonferroni correction was used to correct for multiple testing where significant p -values were multiplied by 6 (i.e. the number of tested SNPs). A two-sided p -value <0.05 was considered statistically significant.

Power calculations were performed using Genetic Power Calculator (available at <http://ibgwww.colorado.edu/~pshaun/gpc/>) [32]. The prevalence of gestational diabetes mellitus was assumed to be 2%. The present study has $> 80\%$ power, under a multiplicative model, to detect an effect size of 1.3 (as measured in terms of genotypic relative risk) when the frequency of the predisposing allele is 15% (for $\alpha=0.05$). The study has at least 80% power to detect a genotypic relative risk of 1.22 (for $\alpha=0.05$) when the predisposing allele frequency is $>30\%$.

Results

Characteristics of the subjects Women with gestational diabetes mellitus were slightly older than pregnant non-diabetic control women (32.3 ± 0.2 vs 30.5 ± 0.1 , $p < 1 \times 10^{-10}$). The genotype and allele frequency distributions of all polymorphisms studied are presented in Table 1.

TCF7L2 rs7903146 The CC, CT and TT genotype frequencies of the *TCF7L2* rs7903146 variant differed significantly between women with gestational diabetes mellitus and control subjects (46.3, 43.6 and 10.1% vs 58.5, 35.3 and 6.2%, $p = 3.7 \times 10^{-6}$, corrected p-value [Pc] for multiple testing $P_c = 2.2 \times 10^{-5}$). The T-allele was associated with an increased risk of gestational diabetes mellitus (OR 1.49 [95% CI 1.28–1.75], $p = 4.9 \times 10^{-7}$ [$P_c = 2.8 \times 10^{-6}$]). Compared with wild-type CC-genotype carriers, heterozygous (CT-genotype) and homozygous (TT-genotype) carriers had a 1.56-fold (95% CI 1.26–1.93, $p = 3.7 \times 10^{-5}$ [$P_c = 0.0002$]) and a 2.05-fold (95% CI 1.41–2.99, $p = 0.0001$ [$P_c = 0.0008$]) increased risk of gestational diabetes mellitus, respectively. Age-adjusted risk of gestational diabetes mellitus for CT-genotype and TT-genotype carriers was 1.60 (95% CI 1.29–1.98, $p = 2 \times 10^{-5}$ [$P_c = 0.0001$]) and 2.08 (95% CI 1.42–3.05, $p = 0.0002$ [$P_c = 0.001$]) respectively. In addition, the effect size was found to change slightly (1.58 [1.27–1.95], $p = 2.8 \times 10^{-5}$ [$P_c = 0.0002$] for CT-genotype; 1.95 [1.33–2.86], $p = 0.0005$ [$P_c = 0.003$] for TT-genotype) when women who were positive for autoantibodies to GAD65 or to protein tyrosine phosphatase or both were removed from analyses. Data on antibody measurements were not available for all subjects.

ADIPOQ +276G>T The T-allele of the +276G>T polymorphism was associated with a slight increased risk of gestational diabetes mellitus (1.17 [1.01–1.36], $p = 0.039$

[Pc=0.23]). In addition, GT-genotype carriers had an increased risk of gestational diabetes mellitus (1.27 [1.04–1.55], p=0.020 [Pc=0.12]) as compared with GG-genotype carriers. A similar effect size (1.26 [1.04–1.53], p=0.018 [Pc=0.11]) was also observed under a dominant model (TT+GT vs GG).

Other polymorphisms The other polymorphisms studied were not significantly associated with gestational diabetes mellitus (*PPARG* Pro12Ala: 1.06 [0.87–1.29], p=0.53; *PPARGC1A* Gly482Ser: 0.96 [0.83–1.10], p=0.54; *FOXO2* -512C>T: 1.01 [0.87–1.16], p=0.94; and *ADRB3* Trp64Arg: 1.22 [0.95–1.56], p=0.12).

Discussion

Gestational diabetes mellitus is a heterogeneous disorder where genetic and environmental factors interact to cause the disease [1, 2]. Studies have demonstrated that gestational diabetes mellitus shares some genetic and phenotypic features with other types of diabetes such as type 1, type 2 and MODY [1, 2]. However, different risk alleles/genes might be operative in different types of diabetes and gestational diabetes mellitus due to the heterogeneous nature of these disorders. During pregnancy beta cells undergo structural and functional changes in response to the increased insulin requirements. These changes include, among others, enhanced glucose-stimulated insulin secretion, increased insulin synthesis and increased beta cell proliferation and islet volume [33]. Genetic factors might therefore predispose to abnormal glucose tolerance during pregnancy by influencing one or more of these physiological processes.

TCF7L2 rs7903146 Our results provide evidence that *TCF7L2* is a major susceptibility gene for gestational diabetes mellitus in Scandinavian women. *TCF7L2*, which maps to chromosome 10q25.3, is a transcription factor belonging to the high mobility group-box transcription factors family [34]. It is involved in the wingless-type MMTV integration site family (WNT) signalling pathway, which is important for the development and growth regulatory mechanisms of the cell [35]. The *TCF7L2* rs7903146 variant was originally associated with type 2 diabetes in individuals from Iceland, Denmark and the USA [12]. This variant was reproducibly associated with type 2 diabetes in subsequent studies [13–19]. The mechanism by which it might lead to deterioration of glucose homeostasis is still unknown. However, recent findings suggest that it is associated with impaired insulin secretion [14, 15, 17]. *TCF7L2* is

abundantly produced in the gut and has been shown to bind to the promoter of proglucagon gene in vitro [36]. Hypothetically, *TCF7L2* might influence insulin secretion through regulation of the insulintropic hormone, glucagon-like peptide-1, which is encoded by the proglucagon gene [37]. A recent report demonstrated that *TCF7L2* is expressed in pancreatic beta cells [13]. Thus, by affecting the WNT signalling, the variant could influence the beta cell proliferation (and hence insulin secretion) that normally takes place during pregnancy [33, 35]. The effect size for gestational diabetes mellitus conferred by this variant is similar to that reported in patients with type 2 diabetes [12]. Also, the allele frequency is comparable to that reported in Scandinavians [12, 15].

ADIPOQ +276G>T *ADIPOQ* is a physiologically active polypeptide hormone derived from adipose tissue with insulin-sensitising properties [38]. Decreased plasma adiponectin during pregnancy has been associated with gestational diabetes mellitus [39]. In addition, reduced *ADIPOQ* mRNA levels in adipose tissue from women with gestational diabetes mellitus have been reported [39]. The common +276G>T variant is one of the most extensively studied variants within the *ADIPOQ* gene. It has been associated with type 2 diabetes, but different populations were found to have different at-risk alleles [20, 21]. Thus, it is possible that another polymorphism in linkage disequilibrium with the +276G>T variant could confer the risk. In the present study, we found a nominal association between the T-allele of the +276G>T variant and gestational diabetes mellitus, but the significance disappeared after correcting for multiple testing. This variant might therefore predispose to gestational diabetes mellitus, but this needs to be further investigated in a larger sample and/or in other populations.

PPARG Pro12Ala *PPARG*, which maps to chromosome 3p25, is a transcription factor with a pivotal role in adipocyte differentiation and function. The Ala allele of the Pro12Ala polymorphism has been consistently associated with reduced risk of type 2 diabetes [22, 40]. In vitro, the Ala allele leads to decreased *PPARG* activity and thereby to decreased transcription of a number of target genes, resulting in increased insulin sensitivity [41]. In the present study, and consistent with our previous finding in a smaller study of Scandinavian and Arabian women [31], we did not find an association between this variant and gestational diabetes mellitus. However, we were unable to rule out a smaller effect size (OR<1.3) of this variant on the risk of gestational diabetes mellitus.

PPARGC1A Gly482Ser *PPARGC1A* is a transcriptional co-activator of several nuclear receptors including *PPARG* and *PPAR* alpha, which plays a role in the transcriptional control of mitochondrial fatty acid beta-oxidation enzymes [42]. *PPARGC1A*, located on chromosome 4p15.1, is expressed in various tissues including adipose tissue, skeletal muscle and pancreas [43]. The common Gly482Ser polymorphism in *PPARGC1A* has been associated with an increased risk of type 2 diabetes [23]. In addition, it influences maximum volume of oxygen uptake (VO₂max) [44] and insulin secretion [45]. We found no effect of this variant on gestational diabetes mellitus in Scandinavian women, a finding which is in agreement with a recent small study in Austrian Europids [46].

FOXC2 -512C>T *FOXC2* is a key regulator of adipocyte metabolism [47]. The common -512C>T polymorphism located in the 5'UTR of the *FOXC2* gene has been associated with enhanced insulin sensitivity and lower plasma triacylglycerols in

female subjects from Scandinavia and in Pima Indian women [24, 48]. This sex-specific association with insulin resistance prompted us to investigate whether this polymorphism also influences the risk of gestational diabetes mellitus. Our results suggest that this polymorphism has no impact on the development of gestational diabetes mellitus in Scandinavian women.

ADRB3 Trp64Arg We and others have previously reported a polymorphism (Trp64Arg) located in the first intracellular loop of the *ADRB3* receptor, which has been associated with early-onset type 2 diabetes, abdominal obesity and features of the metabolic syndrome [49, 50]. Analysis of subsequent studies has shown an association of this polymorphism with features of the metabolic syndrome [40]. In addition, the Trp64Arg polymorphism has been associated with ‘mild’ gestational diabetes mellitus in Austrian [25], but not in Greek [26], Taiwanese [27] or Italian [28] women. However, it was associated with increased weight gain and increased glucose and insulin levels during pregnancy [25, 27]. The present study failed to find a role of this variant in the risk of gestational diabetes mellitus in Scandinavian women.

There are some limitations to the present study. First, we lack detailed phenotypic information on insulin secretion and action during pregnancy. Second, although the study is the largest of its kind, it is not sufficiently powered to detect a weak effect size ($OR < 1.3$). Although insulin resistance is a key feature of gestational diabetes mellitus, genetic variants that affect insulin sensitivity do not seem to increase the risk of gestational diabetes mellitus in Scandinavian women. It could be hypothesised that, due to the massive insulin resistance during pregnancy, we were unable to detect a potential weak effect of the above-mentioned variants on insulin sensitivity.

In conclusion, the *TCF7L2* rs7903146 variant is associated with an increased risk of gestational diabetes mellitus. Given the influence of the variant on insulin secretion [14, 15, 17] and our previous findings of associations of *KCNJ11* E23K, *GCK* – 30G>A and *TCF1* I27L [10, 11] variants with gestational diabetes mellitus, this finding supports the central role of impaired beta cell function in the pathogenesis of gestational diabetes mellitus [3].

Acknowledgements

This work was supported by grants from the Swedish Research Council, the Söderberg foundation, the Lundberg Foundation, the Novo Nordisk Foundation and Diabetes Association in Malmö. We are indebted to M. Svensson, A. Berglund and A. Nilsson for excellent technical assistance.

Duality of interest

The authors declare that no conflict of interest exists in connection with this study.

References

1. Ben-Haroush A, Yogeve Y, Hod M (2004) Epidemiology of gestational diabetes mellitus and its association with Type 2 diabetes. *Diabet Med* 21:103-113
2. Shaat N, Groop L (2007) Genetics of gestational diabetes mellitus. *Curr Med Chem* [in press]
3. Buchanan TA (2001) Pancreatic B-cell defects in gestational diabetes: implications for the pathogenesis and prevention of type 2 diabetes. *J Clin Endocrinol Metab* 86:989-993
4. Poulsen P, Levin K, Petersen I, Christensen K, Beck-Nielsen H, Vaag A (2005) Heritability of insulin secretion, peripheral and hepatic insulin action, and intracellular glucose partitioning in young and old Danish twins. *Diabetes* 54:275-283
5. Solomon CG, Willett WC, Carey VJ et al (1997) A prospective study of pregravid determinants of gestational diabetes mellitus. *JAMA* 278:1078-1083
6. Williams MA, Qiu C, Dempsey JC, Luthy DA (2003) Familial aggregation of type 2 diabetes and chronic hypertension in women with gestational diabetes mellitus. *J Reprod Med* 48:955-962
7. Florez JC, Burt N, de Bakker PI et al (2004) Haplotype structure and genotype-phenotype correlations of the sulfonylurea receptor and the islet ATP-sensitive potassium channel gene region. *Diabetes* 53:1360-1368
8. Rose CS, Ek J, Urhammer SA, Glumer C et al (2005) A -30G>A polymorphism of the {beta}-cell-specific glucokinase promoter associates with hyperglycemia in the general population of whites. *Diabetes* 54:3026-3031
9. Holmkvist J, Cervin C, Lyssenko V et al (2006) Common variants in HNF-1 alpha and risk of type 2 diabetes. *Diabetologia* 49:2882-2891
10. Shaat N, Ekelund M, Lernmark A et al (2005) Association of the E23K polymorphism in the *KCNJ11* gene with gestational diabetes mellitus. *Diabetologia* 48:2544-2551
11. Shaat N, Karlsson E, Lernmark A et al (2006) Common variants in MODY genes increase the risk of gestational diabetes mellitus. *Diabetologia* 49:1545-1551
12. Grant SF, Thorleifsson G, Reynisdottir I et al (2006) Variant of transcription factor 7-like 2 (TCF7L2) gene confers risk of type 2 diabetes. *Nat Genet.* 38:320-323
13. Cauchi S, Meyre D, Dina C et al (2006) Transcription factor TCF7L2 genetic study in the French population: Expression in human {beta}-cells and adipose tissue and strong association with type 2 diabetes. *Diabetes* 55:2903-2908

14. Florez JC, Jablonski KA, Bayley N et al (2006) TCF7L2 polymorphisms and progression to diabetes in the Diabetes Prevention Program. *N Engl J Med* 355:241-250
15. Saxena R, Gianniny L, Burt NP et al (2006) Common single nucleotide polymorphisms in TCF7L2 are reproducibly associated with type 2 diabetes and reduce the insulin response to glucose in non-diabetic individuals. *Diabetes* 55:2890-2895
16. Van Vliet-Ostaptchouk JV, Shiri-Sverdlov R, Zhernakova A et al (2006) Association of variants of transcription factor 7-like 2 (TCF7L2) with susceptibility to type 2 diabetes in the Dutch Breda cohort. *Diabetologia* 50:59-62
17. Damcott CM, Pollin TI, Reinhart LJ et al (2006) Polymorphisms in the transcription factor 7-like 2 (TCF7L2) gene are associated with type 2 diabetes in the Amish: Replication and evidence for a role in both insulin secretion and insulin resistance. *Diabetes* 55:2654-2659
18. Groves CJ, Zeggini E, Minton J et al (2006) Association analysis of 6,736 U.K. subjects provides replication and confirms TCF7L2 as a type 2 diabetes susceptibility gene with a substantial effect on individual risk. *Diabetes* 55:2640-2644
19. Chandak GR, Janipalli CS, Bhaskar S et al (2006) Common variants in the TCF7L2 gene are strongly associated with type 2 diabetes mellitus in the Indian population. *Diabetologia* 50:63-67
20. Hara K, Boutin P, Mori Y et al (2002) Genetic variation in the gene encoding adiponectin is associated with an increased risk of type 2 diabetes in the Japanese population. *Diabetes* 51:536-540
21. Hu FB, Doria A, Li T et al (2004) Genetic variation at the adiponectin locus and risk of type 2 diabetes in women. *Diabetes* 53:209-213
22. Altshuler D, Hirschhorn JN, Klannemark M et al (2000) The common PPARgamma Pro12Ala polymorphism is associated with decreased risk of type 2 diabetes. *Nat Genet* 26:76-80
23. Barroso I, Luan J, Sandhu MS et al (2006) Meta-analysis of the Gly482Ser variant in PPARGC1A in type 2 diabetes and related phenotypes. *Diabetologia* 49:501-505
24. Ridderstrale M, Carlsson E, Klannemark M et al (2002) FOXC2 mRNA expression and a 5' untranslated region polymorphism of the gene are associated with insulin resistance. *Diabetes* 51:3554-3560
25. Festa A, Krugluger W, Shnawa N, Hopmeier P, Haffner SM, Schernthaner G (1999) Trp64Arg polymorphism of the beta3-adrenergic receptor gene in pregnancy: association with mild gestational diabetes mellitus. *J Clin Endocrinol Metab* 84:1695-1699

26. Alevizaki M, Thalassinou L, Grigorakis SI et al (2000) Study of the Trp64Arg polymorphism of the beta3-adrenergic receptor in Greek women with gestational diabetes. *Diabetes Care* 23:1079-1083
27. Tsai PJ, Ho SC, Tsai LP et al (2004) Lack of relationship between beta3-adrenergic receptor gene polymorphism and gestational diabetes mellitus in a Taiwanese population. *Metabolism* 53:1136-1139
28. Fallucca F, Dalfra MG, Sciuillo E et al (2006) Polymorphisms of insulin receptor substrate 1 and beta(3)-adrenergic receptor genes in gestational diabetes and normal pregnancy. *Metabolism* 55:1451-1456
29. Lind T, Phillips PR (1991) Influence of pregnancy on the 75-g OGTT. A prospective multicenter study. The Diabetic Pregnancy Study Group of the European Association for the Study of Diabetes. *Diabetes* 40 (Suppl 2):8-13
30. Lernmark B, Elding-Larsson H, Hansson G, Lindberg B, Lynch K, Sjoblad S (2004) Parent responses to participation in genetic screening for diabetes risk. *Pediatr Diabetes* 5:174-181
31. Shaat N, Ekelund M, Lernmark A et al (2004) Genotypic and phenotypic differences between Arabian and Scandinavian women with gestational diabetes mellitus. *Diabetologia* 47:878-884
32. Purcell S, Cherny SS, Sham PC (2003) Genetic Power Calculator: design of linkage and association genetic mapping studies of complex traits. *Bioinformatics* 19:149-150. Available from <http://ibgwww.colorado.edu/~pshaun/gpc/> last accessed 09 January 2007
33. Sorenson RL, Brelje TC (1997) Adaptation of islets of Langerhans to pregnancy: beta-cell growth, enhanced insulin secretion and the role of lactogenic hormones. *Horm Metab Res* 29:301-307
34. Duval A, Busson-Leconiat M, Berger R, Hamelin R (2000) Assignment of the TCF-4 gene (TCF7L2) to human chromosome band 10q25.3. *Cytogenet Cell Genet* 88:264-265
35. Douglas KR, Brinkmeier ML, Kennell JA et al (2001) Identification of members of the Wnt signaling pathway in the embryonic pituitary gland. *Mamm Genome* 12:843-851
36. Yi F, Brubaker PL, Jin T (2005) TCF-4 mediates cell type-specific regulation of proglucagon gene expression by beta-catenin and glycogen synthase kinase-3beta. *J Biol Chem* 280:1457-1464
37. Meier JJ, Nauck MA (2005) Glucagon-like peptide 1 (GLP-1) in biology and pathology. *Diabetes Metab Res Rev* 21:91-117

38. Diez JJ, Iglesias P (2003) The role of the novel adipocyte-derived hormone adiponectin in human disease. *Eur J Endocrinol* 148:293-300
39. Ranheim T, Haugen F, Staff AC, Braekke K, Harsem NK, Drevon CA (2004) Adiponectin is reduced in gestational diabetes mellitus in normal weight women. *Acta Obstet Gynecol Scand* 83:341-347
40. Parikh H, Groop L (2004) Candidate genes for type 2 diabetes. *Rev Endocr Metab Disord* 5:151-176
41. Deeb SS, Fajas L, Nemoto M et al (1998) A Pro12Ala substitution in PPARgamma2 associated with decreased receptor activity, lower body mass index and improved insulin sensitivity. *Nat Genet* 20:284-287
42. Vega RB, Huss JM, Kelly DP (2000) The coactivator PGC-1 cooperates with peroxisome proliferator-activated receptor alpha in transcriptional control of nuclear genes encoding mitochondrial fatty acid oxidation enzymes. *Mol Cell Biol* 20:1868-1876
43. Esterbauer H, Oberkofler H, Krempler F, Patsch W (1999) Human peroxisome proliferator activated receptor gamma coactivator 1 (PPARGC1) gene: cDNA sequence, genomic organization, chromosomal localization, and tissue expression. *Genomics* 62:98-102
44. Ling C, Poulsen P, Carlsson E et al (2004) Multiple environmental and genetic factors influence skeletal muscle PGC-1alpha and PGC-1beta gene expression in twins. *J Clin Invest* 114:1518-1526
45. Muller YL, Bogardus C, Pedersen O, Baier L (2003) A Gly482Ser missense mutation in the peroxisome proliferator-activated receptor gamma coactivator-1 is associated with altered lipid oxidation and early insulin secretion in Pima Indians. *Diabetes* 52:895-898
46. Leipold H, Knoefler M, Gruber C, Huber A, Haslinger P, Worda C (2006) Peroxisome proliferator-activated receptor gamma coactivator-1alpha gene variations are not associated with gestational diabetes mellitus. *J Soc Gynecol Investig* 13:104-107
47. Cederberg A, Gronning LM, Ahren B, Tasken K, Carlsson P, Enerback S (2001) FOXC2 is a winged helix gene that counteracts obesity, hypertriglyceridemia, and diet-induced insulin resistance. *Cell* 106:563-573
48. Kovacs P, Lehn-Stefan A, Stumvoll M, Bogardus C, Baier LJ (2003) Genetic variation in the human winged helix/forkhead transcription factor gene FOXC2 in Pima Indians. *Diabetes* 52:1292-1295

49. Widen E, Lehto M, Kanninen T, Walston J, Shuldiner AR, Groop LC (1995) Association of a polymorphism in the beta 3-adrenergic-receptor gene with features of the insulin resistance syndrome in Finns. *N Engl J Med* 333:348-351
50. Walston J, Silver K, Bogardus C et al (1995) Time of onset of non-insulin-dependent diabetes mellitus and genetic variation in the beta 3-adrenergic-receptor gene. *N Engl J Med* 333:343-347

Table 1. Genotype and allele distributions and corresponding odds ratios for GDM

Polymorphism (rs number)	Genotype or allele	GDM, n (%)	Controls subjects, n (%)	GDM, allelic effect, additive models ^a	GDM, recessive model ^a	GDM, dominant model ^a
<i>TCF7L2</i> IVS3C>T (rs7903146)	CC	271 (46.3)	650 (58.5)	1		
	CT	255 (43.6)	392 (35.3)	1.56 (1.26–1.93) ^c		
	TT	59 (10.1)	69 (6.2) ^b	2.05 (1.41–2.99) ^d	1.69 (1.18–2.44) ^f	1.63 (1.34–2.0) ^g
	T	373 (31.9)	530 (23.8)	1.49 (1.28–1.75) ^c		
<i>ADIPOQ</i> +276G>T (rs1501299)	GG	301 (46.9)	646 (52.7)	1		
	GT	285 (44.5)	482 (39.4)	1.27 (1.04–1.55)		
	TT	55 (8.6)	97 (7.9)	1.22 (0.85–1.74)	1.09 (0.77–1.54)	1.26 (1.04–1.53)
	T	395 (30.8)	676 (27.6)	1.17(1.01–1.36) ^h		
<i>PPARG</i> Pro12Ala (rs1801282)	Pro/Pro	468 (73.5)	918 (74.5)	1		
	Pro/Ala	158 (24.8)	298 (24.2)	1.04 (0.83–1.30)		
	Ala/Ala	11 (1.7)	16 (1.3)	1.35 (0.62–2.93)	1.34 (0.62–2.89)	1.06 (0.85–1.31)
	Ala	180 (14.1)	330 (13.4)	1.06 (0.87–1.29)		
<i>PPARGC1A</i> Gly482Ser (rs8192678)	Gly/Gly	284 (44.1)	533 (43.5)	1		
	Gly/Ser	294 (45.7)	548 (44.8)	1.01 (0.82–1.23)		
	Ser/Ser	66 (10.2)	143 (11.7)	0.87 (0.63–1.20)	0.86 (0.63–1.18)	0.98 (0.81–1.19)
	Ser	426 (33.1)	834 (34.1)	0.96 (0.83–1.10)		
<i>FOXC2</i> -512C>T	TT	244 (38.4)	456 (37.9)	1		
	CT	291 (45.8)	568 (47.2)	0.96 (0.78–1.18)		
	CC	100 (15.8)	180 (14.9)	1.04 (0.78–1.39)	1.06 (0.82–1.39)	0.98 (0.80–1.19)
	C	491 (38.7)	928 (38.5)	1.01 (0.87–1.16)		
<i>ADRB3</i> Trp64Arg (rs4994)	Trp/Trp	534 (83.6)	1060(86.4)	1		
	Trp/Arg	100 (15.6)	158 (12.9)	1.26 (0.96–1.65)		
	Arg/Arg	5 (0.8)	9 (0.7)	1.10 (0.37–3.31)	1.07 (0.36–3.20)	1.25 (0.96–1.63)
	Arg	110 (8.6)	176 (7.2)	1.22 (0.95–1.56)		

GDM, gestational diabetes mellitus

^a OR (95% CI)

^b $p=3.7 \times 10^{-6}$ ($P_c=2.2 \times 10^{-5}$) for difference in genotype frequencies between women with and without GDM

^c $p=3.7 \times 10^{-5}$ ($P_c=0.0002$) for comparison of CT vs CC between women with and without GDM

^d $p=0.0001$ ($P_c=0.0008$) for comparison of TT vs CC between women with and without GDM

^e $p=4.9 \times 10^{-7}$ ($P_c=2.8 \times 10^{-6}$) for difference in T-allele frequencies between women with and without GDM

^f $p=0.004$ ($P_c=0.02$) for comparison of TT vs CT/CC (recessive model) between women with and without GDM

^g $p=1.7 \times 10^{-6}$ ($P_c=1 \times 10^{-5}$) for comparison of TT/CT vs CC (dominant model) between women with and without GDM

^h $p=0.039$ ($P_c=0.23$) for difference in T-allele frequencies between women with and without GDM

Electronic supplementary material

ESM Table 1 Primers and probes used for TaqMan allelic discrimination assay

Polymorphism (rs number)	Forward primer (5'→3')	Reverse primer (5'→3')	Probe (5'FAM)	Probe (5'VIC)
<i>TCF7L2</i> IVS3C>T (rs7903146)	CCTCAAAACCTAGC ACAGCTGTTAT	TGAAAACCTAAGG GTGCCTCATACG	CTAAGCACTTTTT AGATACTATAT	TAAGCACTTTTTA GATATTATAT
<i>ADIPOQ</i> +276G>T (rs1501299)	TTCATCACAGACCT CCTACACTGA	TCCCTGTGTCTAG GCCTTAGTTAAT	AACTATATGAAG TCATTCAT	ACTATATGAAGGC A TTCAT
<i>PPARG</i> Pro12Ala (rs1801282)	GTTATGGGTGAAAC TCTGGGAGATT	GCAGACAGTGTAT CAGTGAAGGAAT	CTATTGACGCAGA AAG	CTCCTATTGACCC AGAAAG
<i>PPARGC1A</i> Gly482Ser (rs8192678)	TGGAGAATTGTTCA TTACTGAAATCACT GT	GGTCATCCCAGTC AAGCTGTTTT	ACAAGACCAGTG AACTG	CAAGACCGGTGA ACTG
<i>FOXC2</i> -512C>T	CGGGTGATTGGCTC AAAGTT	GCCAAGTCCCTTT TAGGGATTG	TCGCTTTCAGCAA G AAGATTTTTGAAA CT-(BHQ1)	(TAM)- TCGCTTTCAGCAA GAAGACTTTTGAA ACT-(BHQ2)
<i>ADRB3</i> Trp64Arg (rs4994)	GTTGGTCATGGTCT GGAGTCT	GCAACCTGCTGGT CATCGT	ATCGCCCGGACTC	CATCGCCTGGACT C

ESM Table 2 Primers used for template PCRs

Polymorphism (rs number)	Forward primer (5'→3')	Reverse primer (5'→3')	Fragment size (bp)	Annealing temperature (°C)
<i>TCF7L2</i> IVS3C>T (rs7903146)	GGAGAAAGCAGGATTG AGCAG	TGCCTTCCCTGTAACTG TGGT	251	64
<i>ADIPOQ</i> +276G>T (rs1501299)	AGAAAGCAGCTCCTAG AAGT	GGCACCATCTACACTC ATCC	518	58
<i>PPARG</i> Pro12Ala (rs1801282)	CAAACCCCTATTCCAT GCTG	CCTTACATAAATGCCC CCAC	157	59
<i>PPARGC1A</i> Gly482Ser (rs8192678)	GGGGTCTTTGAGAAAA TAAGG	CAAGTCCTCAGTCCTC AC	611	58
<i>FOXC2</i> -512C>T	GTCTTAGAGCCGACGG ATTCCTG	TGGGGACCAAGGTGGA CCCTCG	306	63
<i>ADRB3</i> Trp64Arg (rs4994)	CGCCCAATACCGCCAA CACC	CCACCAGGAGTCCCAT CACC	210	63

ESM Table 3 PCR reagents used for amplification of template PCRs ^a

Polymorphism (rs number)	PCR buffer	dNTP (nmol) ^d	Primers (pmol)	MgCl ₂ (nmol)	Betaine (μmol) ^e	DMSO (%)
<i>TCF7L2</i> IVS3C>T (rs7903146)	1x Pharmacia Amersham ^b	8	20	–	20	–
<i>ADIPOQ</i> +276G>T (rs1501299)	1x (NH ₄) ₂ SO ₄ ^c	4	20	60	20	–
<i>PPARG</i> Pro12Ala (rs1801282)	1x Pharmacia Amersham	4	20	–	20	–
<i>PPARGC1A</i> Gly482Ser (rs8192678)	1x Pharmacia Amersham	8	20	30	10	5
<i>FOXC2</i> -512C>T	1x (NH ₄) ₂ SO ₄	4	20	60	20	5
<i>ADRB3</i> Trp64Arg (rs4994)	1x (NH ₄) ₂ SO ₄	8	20	120	20	–

^a PCR amplification was carried out with 3 mm of dried blood spots in a total volume of 40 μl with 1.5–2.5 U of Taq polymerase (New England Biolabs, Beverly, MA, USA)

^b Produced by Amersham Pharmacia Biotech, Uppsala, Sweden

^c (NH₄)₂SO₄ buffer contains: 16 mmol/l (NH₄)₂SO₄, 67 mmol/l Tris (pH 8.8) and 0.01% Tween 20

^d Produced by Fermentas, St Leon-Rot, Germany

^e Produced by Sigma-Aldrich Sweden, Stockholm, Sweden