



LUND UNIVERSITY

Is there an association between female circumcision and perinatal death?

Essén, Birgitta; Bodker, Birgit; Sjöberg, Nils-Otto; Gudmundsson, Saemundur; Östergren, Per-Olof; Langhoff-Roos, Jens

Published in:
Bulletin of the World Health Organization

2002

[Link to publication](#)

Citation for published version (APA):

Essén, B., Bodker, B., Sjöberg, N.-O., Gudmundsson, S., Östergren, P.-O., & Langhoff-Roos, J. (2002). Is there an association between female circumcision and perinatal death? *Bulletin of the World Health Organization*, 80(8), 629-632.

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=12219153&dopt=Abstract

Total number of authors:
6

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Is there an association between female circumcision and perinatal death?

Birgitta Essén,¹ Birgit Bødker,² N-O. Sjöberg,³ Saemundur Gudmundsson,⁴ P-O. Östergren,⁵
& Jens Langhoff-Roos⁶

Objective In Sweden, a country with high standards of obstetric care, the high rate of perinatal mortality among children of immigrant women from the Horn of Africa raises the question of whether there is an association between female circumcision and perinatal death.

Method To investigate this, we examined a cohort of 63 perinatal deaths of infants born in Sweden over the period 1990–96 to circumcised women.

Findings We found no evidence that female circumcision was related to perinatal death. Obstructed or prolonged labour, caused by scar tissue from circumcision, was not found to have any impact on the number of perinatal deaths.

Conclusion The results do not support previous conclusions that genital circumcision is related to perinatal death, regardless of other circumstances, and suggest that other, suboptimal factors contribute to perinatal death among circumcised migrant women.

Keywords Circumcision, Female/adverse effects; Labor complications/etiology; Infant mortality; Transients and migrants; Cohort studies; Africa, Eastern; Sweden (*source: MeSH, NLM*).

Mots clés Infibulation/effets indésirables; Accouchement compliqué/étiologie; Mortalité nourrisson; Population passage et immigrants; Etude cohorte; Afrique orientale; Suède (*source: MeSH, INSERM*).

Palabras clave Circuncisión femenina/efectos adversos; Complicaciones del trabajo de parto/etiología; Mortalidad infantil; Transeúntes y migrantes; Estudios de cohortes; África Oriental; Suecia (*fuentes: DeCS, BIREME*).

Bulletin of the World Health Organization 2002;80:629-632.

Voir page 631 le résumé en français. En la página 632 figura un resumen en español.

Introduction

Much has been written about complications arising from female circumcision (defined by WHO as female genital mutilation involving the partial or total removal of external genitalia, whether for cultural or other nontherapeutic reasons) in Africa (1, p. 23–36). Although it has been claimed that scar or soft-tissue dystocia from circumcision can prolong the second stage of labour and lead to cerebral damage, asphyxia, and fetal death, few controlled studies have been published (2–4). Also, little attention has been paid to obstetric complications arising from genital mutilation among migrant women who move from a low-resource to a high-resource country (5–11).

A recent investigation showed that perinatal mortality among women who emigrated to Sweden from sub-Saharan Africa (predominantly Somalia) was greater than that of their Swedish counterparts, even after adjustments were made for background factors and perinatal risk factors (12). In three interview studies of circumcised Somali women who had immigrated to high-income countries (the USA and Sweden), the women expressed concern that Western health care providers would not know how to deal with circumcised women, especially during delivery (13–15). Somali women in

Sweden perceived Caesarean section as a life-threatening event and many reduced their food intake to prevent dystocia and to avoid operative deliveries. A similar fear was expressed by Somali women in the USA, and they favoured obstetricians with low rates of Caesarean sections. However, the Somali women in Sweden did not associate circumcision with obstetric complications, such as obstructed labour or perinatal death.

The present study was designed to test the hypothesis that genital circumcision is a contributing factor to the increased rate of perinatal death among infants of immigrant women who gave birth in a community with a high standard of obstetric care.

Materials and methods

A perinatal audit of medical records in the national Swedish Medical Birth Register was made for all 63 perinatal deaths (defined as stillbirth after 28 weeks of gestation or death within the first week of life) that occurred in Sweden over the period 1990–96. The records represented the full cohort of pregnant women from the Horn of Africa who lived in Sweden during this period ($n = 32$ for Ethiopia/Eritrea; $n = 31$ for Somalia).

¹ Department of Obstetrics and Gynaecology, University Hospital MAS, Lund University, S-205 02 Malmö, Sweden (email: Birgitta.Essen@obst.mas.lu.se). Correspondence should be addressed to this author.

² Senior Lecturer, Department of Obstetrics and Gynaecology, Hillerød Hospital, Copenhagen, Denmark.

³ Professor, Department of Obstetrics and Gynaecology, University Hospital MAS, Lund University, Malmö, Sweden.

⁴ Senior Lecturer, Department of Obstetrics and Gynaecology, University Hospital MAS, Lund University, Malmö, Sweden.

⁵ Senior Lecturer, Department of Community Medicine, University Hospital MAS, Lund University, Malmö, Sweden.

⁶ Senior Lecturer, Department of Obstetrics and Gynaecology, and Department of International Health, University of Copenhagen, Copenhagen, Denmark.

Narratives based on individual records were obtained for each birth and contained in-depth descriptions of relevant time-related events, including medical history, social conditions, communication, surveillance, interventions and outcome. The narratives were limited to data obtained from the records and noted whether pertinent information was missing. All of the women had been circumcised, although the exact form of vulva trauma was not always described in detail. The commonest form of genital cutting among girls from the Horn of Africa is infibulation (16), an extensive form of mutilation involving en-bloc removal of the clitoris, minor labia, parts of the major labia and stitching together the remains of the major labia to leave a small opening at the lower part of the vulva.

An expert panel of three obstetricians and one neonatologist was convened to review the 63 case narratives and identify suboptimal factors that were likely to have contributed to perinatal death. Criteria for identifying suboptimal factors were primarily evidence based and adopted from the EuroNatal study of perinatal mortality in Europe (17). The study was approved by the Ethics Committee, University of Lund, Sweden.

Results

Background and risk factors for the 63 circumcised mothers and their infants are shown in Table 1. Caesarean section was performed on 16% of the women, with threatening fetal distress being the commonest indication for performing it, but no indication of disproportion, shoulder dystocia or obstructed labour was found.

Table 2 presents a summary of the audits for the perinatal deaths. No documented evidence has been found that antenatal death is considered to be related to circumcision (7 severe malformations, 16 intrauterine growth retarded, 4 placental abruption, 1 twin transfusion syndrome, 3 stillbirth with positive culture of maternal cervical streptococci group B and 8 unexplained). There were eight intrapartum deaths: four were placental abruptions and four were due to fetal distress. However, none of the intrapartum deaths was related to genital mutilation and obstructed labour based on the audit of the medical records. Of the neonatal deaths, one involved a nullipara with a prolonged first stage of labour (25 hours). Retrospective analysis indicated that cardiotocography (CTG) showed signs of severe fetal distress for three hours without appropriate action being taken. Nevertheless, the infant was delivered vaginally without difficulty but died due to complications from neonatal distress. In general, misinterpretation of CTGs by midwives or obstetricians and mothers who avoided Caesarean section were important suboptimal factors that probably contributed to fatal outcomes in the cohort.

Discussion

None of the perinatal deaths in the present study was considered to be related to complications caused by circumcision of the mothers. Neither obstructed labour nor prolonged second-stage of labour was common and had no impact on the number of perinatal deaths. In contrast, perinatal deaths among the study women were associated with suboptimal factors, such as maternal behaviour (mother avoiding operative delivery), verbal miscommunication and insufficient obstetric care, including misinterpretation of intrapartum CTG.

Table 1. Background and risk factors for 63 circumcised study mothers, Sweden, 1990–96

Background and risk factors	No. of mothers
Maternal age 19–40 years	62 (98.4) ^a
Smokers in early pregnancy	1 (1.6)
Antenatal care since the first trimester ^b	33 (65.0)
No ultrasound screening in the second trimester	11 (17.4)
Nullipara	21 (33.3)
Parity >4	4 (6.4)
Gestational age (weeks)	
<28	6 (9.5)
28–33	21 (33.3)
>33	36 (57.1)
Twin pregnancy	1 (1.6)
Maternal disease ^c	17 (27.0)
Caesarean section	10 (15.9)
SGA ^d	25 (39.7)
Neonatal distress ^e	16 (25)
Severe malformation	7 (11)

^a Figures in parentheses are percentages.

^b $n=51$.

^c HIV-infection, diabetes, epilepsy or cardiac disease.

^d Small for gestational age (two data points missing).

^e Apgar score 5' <7 ($n=23$).

Our results agree with those of a recent publication that found no relation between obstructed labour due to tough scars around the vaginal exit, including those from female genital mutilation, since the elasticity of the birth canal is not affected (1, p. 42). The results suggest that the elasticity of the birth canal is no more affected by circumcision scar tissue than by scar tissue induced by episiotomy.

For infibulated women there is a major risk of severe perineum tearing unless defibulation is performed (10). Although circumcision might not cause obstructed labour, it is important that defibulation be performed and that the women give birth in hospital. In the present study, defibulation or episiotomy were always performed when needed, which may account for the lack of correlation between circumcision and obstructed labour (8). No evidence of vaginal stenosis was reported, in contrast to the findings of earlier studies of non-nulliparous circumcised women in Somalia (17, 18).

High birth weight is known to be associated with obstructed labour. A total of 40% of infants born to circumcised women were small for gestational age (birth weight > 2 standard deviations below the mean) (19). Theoretically, fetal growth retardation might lead to easier vaginal delivery, with less risk of perineal injury and dystocia. On the other hand, it can be a cause for emergency interventions due to imminent asphyxia. Dystocia was not described in any of the cases in the present study.

Fetal distress was noted in some cases and was related to perinatal death because the mother refused emergency Caesarean section (Table 2). Because of fear of delivery complications and maternal death, the mothers often delayed or refused Caesarean section, even when experiencing severe symptoms such as vaginal hemorrhage or if there was a risk of fetal distress (13). The women's experiences in their countries of origin, where maternal mortality rates are high, resulted in

Table 2. Audit of 63 perinatal deaths to circumcised study mothers, Sweden, 1990–96

No. of deaths and relation to delivery	Reason for perinatal death	Comments	Circumcision related
Antenatal 39	Stillbirths	Malformation or IUGR ^a	Not likely
Intrapartal 4	Placental abruption and fetal distress	2 mothers refused C/S, ^b 1 case of misinterpreted CTG ^c	Not likely
4	Fetal distress ^d	1 mother refusing C/S, 2 cases of misinterpreted CTG	Not likely
Neonatal 4	Severe malformation		Not likely
1	Sudden infant death		Not likely
3	IRDS, ^e prematurely		Not likely
1	IRDS and sepsis		Not likely
2	IRDS and neonatal distress ^f	2 mothers refusing C/S	Not likely
5	Neonatal distress	3 cases of misinterpreted CTG	Not likely

^a Intrauterine growth retardation.^b Emergency Caesarean section.^c Cardiotocography.^d Pathological intrapartal CTG.^e Idiopathic Respiratory Distress Syndrome.^f Apgar score 5' < 7.

maternal behaviours and pregnancy strategies that the Swedish caregivers had not considered. These factors, combined with miscommunication because of a lack of interpreters and suboptimal medical care (e.g. misinterpretation of CTG signs of fetal distress), heightened the risk of perinatal mortality. The high observed incidence of intrapartal or neonatal death was thus not related to maternal circumcision in this study.

It could be asked whether the women in our study were representative of their ethnic groups, since they might have represented a positively selected group with long-term complications due to circumcision (such as keloid scarring or cysts) (16). Although we found no evidence that ethnicity per se was an independent risk factor for prolonged labour (20–22), it was not possible to control the studies with non-circumcised immigrant women from the Horn of Africa.

Our findings should not be seen as a reason to vindicate the practice of female circumcision. On the contrary, it might help to focus issues concerning circumcision and underscore the need for intervention, since for the majority of circumcised women the consequences of giving birth might be totally different from those in the present study. Also, in developing countries with poor obstetric care it is difficult to obtain reliable data on birth labour. If the notion that circumcision is an important cause of obstetric risk among immigrant women is unwarranted, it might lead to

attention being focused on the wrong issue. This could result in inadequate intervention measures, rather than enhancing correct strategies against obstetric risk. An increase in the familiarity of Swedish obstetricians with these issues may decrease the perinatal mortality among immigrant women from the Horn of Africa by focusing on patient education, interpersonal communication, and improved fetal surveillance.

The UN recommends that governments adopt clear national policies to abolish female genital mutilation and intensify educational programmes about its harmfulness (23). Prevention of such mutilation is a matter of not only educating families but also health care providers by offering them correct information. In this way, the providers could better focus their work and give the best care to circumcised women in both low- and high-resource areas. ■

Acknowledgements

We thank Dr Jan Hendrik Richardus and Professor Gorm Greisen for their cooperation. This study was supported by grants from the Faculty of Medicine, University of Lund; the University Hospital MAS, Malmö; and the Samaritan and Mayflower Foundations, Sweden.

Conflicts of interest: none declared.

Résumé

Existe-t-il une association entre la circoncision féminine et la mortalité périnatale ?

En Suède, pays ayant des normes élevées en matière de soins obstétricaux, l'importance de la mortalité périnatale chez les enfants de femmes immigrées originaires de la Corne de l'Afrique soulève la question d'une association éventuelle entre la circoncision féminine et les décès périnatals. Dans cette optique, nous avons examiné une cohorte de 63 décès périnatals survenus chez des enfants nés en Suède en 1990-1996 de mères circoncises, et n'avons rien trouvé qui indique une relation entre la circoncision féminine et le décès

périnatal. Il n'a pas été trouvé qu'une dystocie mécanique ou un travail prolongé, dus à la présence de tissu cicatriciel résultant de la circoncision, ait un impact sur le nombre de décès périnatals. Les résultats ne confirment pas ceux de travaux antérieurs faisant état d'une relation entre la circoncision féminine et les décès périnatals, indépendamment des autres circonstances, et laissent à penser que d'autres facteurs, suboptimaux, contribuent à la mortalité périnatale chez les enfants nés de femmes immigrées circoncises.

Resumen

¿Hay relación entre la mutilación genital femenina y la mortalidad perinatal?

En Suecia, país que cuenta con una atención obstétrica de alto nivel, la alta tasa de mortalidad perinatal observada entre los hijos de las mujeres inmigrantes del Cuerno de África obliga a preguntarse si no existirá una relación entre la mutilación genital femenina y la mortalidad perinatal. A fin de investigar esa posibilidad, examinamos una cohorte de 63 defunciones perinatales de lactantes dados a luz en Suecia durante el período 1990-1996 por mujeres sometidas a ese tipo de mutilación. No se

observó efecto alguno del parto obstruido o prolongado, consecuencia del tejido cicatrizal provocado por la mutilación, en el número de defunciones perinatales. Los resultados no respaldan conclusiones anteriores que apuntaban que la mutilación genital guarda relación con la mortalidad perinatal, con independencia de otras circunstancias, y lleva a pensar que hay otros factores subóptimos que contribuyen a las defunciones perinatales entre las mujeres migrantes mutiladas.

References

1. *Female genital mutilation. An overview*. Geneva: World Health Organization; 1998.
2. De Silva S. Obstetric sequel of female circumcision. *European Journal of Obstetrics, Gynecology and Reproductive Biology* 1989;32:233-40.
3. Press F, Katz M, Leiberman JR, Shoham I, Glezerman M. Obstetric performance in Ethiopian immigrants compared with Israeli parturients. *Israeli Journal of Medical Science* 1993;29:403-7.
4. Berardi JC, Teillet JF, Laloux V, Aillane P, Franjou MH. [Obstetric consequences of female circumcision. Study in 71 circumcised African women]. *Journal of Gynecology, Obstetrics and Biological Reproduction (Paris)* 1985;14:743-6 (in French).
5. Arbesman M, Kahler L, Buck GM. Assessment of the impact of female circumcision on the gynecological, genitourinary, and obstetrical health problems of women from Somalia: literature review and case series. *Women Health* 1993;20(3):27-42.
6. Baker CA, Gilson GJ, Vill MD, Curet LB. Female circumcision: obstetric issues. *American Journal of Obstetrics and Gynecology* 1993;169(6):1616-8.
7. Elchalal U, Ben-Ami B, Gillis R, Brzenzinski A. Ritualistic female genital mutilation: current status and future outlook. *Obstetric and Gynecological Surveys* 1997;52(10):643-51.
8. Omar M. Female circumcision and child mortality in urban Somalia. *Genus* 1991;47(3): 203-23.
9. Lightfoot-Klein H, Shaw E. Special needs of ritually circumcised women patients. *Journal of Obstetrics, Gynecology and Neonatal Nursing* 1991;20(2):102-7.
10. Hosken F. *The Hosken report: genital and sexual mutilation of females*. 4th ed. Lexington (MA): WIN News; 1993.
11. Toubia N. Female circumcision as a public health issue. *New England Journal of Medicine* 1994;331:712-6.
12. Essén B, Hanson BS, Ostergren PO, Lindquist PG, Gudmundsson S. Increased perinatal mortality among sub-Saharan immigrants in a city population in Sweden. *Acta Obstetrica Gynecologica et Scandinavica* 2000;79:737-43.
13. Essén B, Johnsdotter S, Hovelius B, Gudmundsson S, Sjöberg NO, Friedman J, et al. Qualitative study of pregnancy and childbirth experiences in Somali women resident in Sweden. *British Journal of Obstetrics and Gynaecology* 2000;107:1507-12.
14. Shaw E. Female circumcision: What kind of maternity care do circumcised women need and can United States caregivers provide it? *American Journal of Nursing* 1985;85:684-7.
15. Beine K, Fullerton J, Palinkas L, Anders B. Conceptions of prenatal care among Somali women in San Diego. *Journal of Nurse-Midwifery* 1995;40:376-81.
16. Dirie MA, Lindmark G. The risk of medical complications after female circumcision. *East African Medical Journal* 1992;69:479-82.
17. Richardus JH, Graafmans WC, van der Pal-de Bruin KM, Amelink-Verburg MP, Verloove-Vanhorick SP, Mackenbach JP. A European concerted action investigating the validity of perinatal mortality as an outcome indicator for the quality of antenatal and perinatal care. *Journal of Perinatal Medicine* 1997;25:313-24.
18. Medical complications of female circumcision: Some preliminary findings. Proceedings of the First National Congress in Health Research, Mogadishu, Somalia, 27-29 January 1986. p.157-163. Faduma Haji Muhammed, Banaadir Hospital, SOMAC, Somalia.
19. Marsal K, Persson PH, Larsen T, Lilja H, Selbing A, Sultan B. Intrauterine growth curves based on ultrasonically estimated foetal weights. *Acta Paediatrica* 1996;85:843-8.
20. Albers LL, Schiff M, Gorwoda JG. The length of active labour in normal pregnancies. *Obstetrics and Gynecology* 1996;87:355-9.
21. Melmed H, Evans MI. Patterns of labour in native and immigrant population in Israel. *Israeli Journal of Medical Science* 1976;12:1405-9.
22. Duignan NM, Studd JWW, Hughes AO. Characteristics of normal labour in different racial groups. *British Journal of Obstetrics and Gynaecology* 1975;82:593.
23. *Female genital mutilation. A joint WHO/UNICEF/UNFPA statement*. Geneva: World Health Organization; 1997.