



LUND UNIVERSITY

Health Effects in Healthy Volunteers in Controlled Experimental Exposure to Diesel Exhaust and Traffic Noise

Gudmundsson, Anders; Xu, Yiyi; Wierzbicka, Aneta; Andersson, Ulla B K; Assarsson, Eva; Axmon, Anna; Barregård, Lars; Berglund, Margareta; Bohgard, Mats; Broberg Palmgren, Karin; Brunskog, Jonas; Gunnskog, Anna-Therese; Hagerman, Inger; Jönsson, Bo A; Nielsen, Jörn; Kåredal, Monica; Nilsson, Patrik; Österberg, Kai; Pagels, Joakim; Poulsen, Torben; Rissler, Jenny; Stockfelt, Leo; Sällsten, Gerd; Albin, Maria

Published in:
Arbete och hälsa

2011

[Link to publication](#)

Citation for published version (APA):

Gudmundsson, A., Xu, Y., Wierzbicka, A., Andersson, U. B. K., Assarsson, E., Axmon, A., Barregård, L., Berglund, M., Bohgard, M., Broberg Palmgren, K., Brunskog, J., Gunnskog, A.-T., Hagerman, I., Jönsson, B. A., Nielsen, J., Kåredal, M., Nilsson, P., Österberg, K., Pagels, J., ... Albin, M. (2011). Health Effects in Healthy Volunteers in Controlled Experimental Exposure to Diesel Exhaust and Traffic Noise. *Arbete och hälsa*, 45(5), 30-30.

Total number of authors:

24

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

30) Health effects in healthy volunteers in controlled experimental exposure to diesel exhaust and traffic noise.

Anders Gudmundsson¹, Yiyi Xu², Aneta Wierzbicka¹, Ulla B.K. Andersson², Eva Assarsson², Anna Axmon², Lars Barregård³, Margareta Berglund³, Mats Bohgard¹, Karin Broberg², Jonas Brunskog⁴, Anna-Therese Gunnskog², Inger Hagerman³, Bo A. Jönsson², Jörn Nielsen², Monica Kåredal², Patrik Nilsson¹, Kai Österberg², Joakim Pagels¹, Torben Poulsen⁴, Jenny Rissler¹, Leo Stockfelt², Gerd Sällsten⁵, and Maria Albin²

¹Ergonomics and Aerosol Technology, Lund University, Sweden

²Occupational and Environmental Medicine, Lund University, Sweden

³Department of Cardiology, Karolinska University Hospital, Huddinge, Sweden

⁴Acoustic Technology and Hearing Systems, Technical University of Denmark, Denmark

⁵Occupational and Environmental Medicine, University of Gothenburg, Sweden

Key-words: diesel exhaust, lung function, acute effects

Introduction: Exposure to diesel exhaust (DE) and traffic noise is common. Previous studies found an inflammatory response in healthy volunteers, but no effect on lung function after one hour diesel exposure (particle mass concentration 350 µg/m³ and NO₂ concentration of 0.62 ppm).

Methods: Eighteen healthy volunteers were exposed in a stainless steel chamber four times, each 3 hours: 1) zero exposure, 2) high DE (300 µg/m³ and NO₂ 1.4 ppm) and low traffic noise, 3) low DE and high traffic noise 75 dB(A) and 4) high DE and high traffic noise.

Immediately before and after exposure we performed medical examination, spirometry, rhinometry and blood sampling (repeated next morning). Symptom scores and peak expiratory flow (PEF) and ECG measurements were assessed before, 15 min, 75 min and 135 min into exposure and after. Generalized Estimating Equation (GEE) model was used to analyze collected data.

Results: Self-rated irritation of the eyes and throat was higher during DE than non-diesel exposure (non-DE), with a statistically significant difference after 75 (eyes) and 135 min (eyes, throat). Signs of irritation in the upper airways were significantly more common after DE (OR=3.2) and tended to be so also for the eyes (OR=3.1, p=0.06). PEF increased during non-diesel exposure, but decreased during DE, with a statistically significant difference after 75 (+4.08 vs -9.58 l/s), and 135 min (+8.11 l/s vs -3.5 l/s). Leukocyte concentrations were higher after exposure to DE than non-DE, and a tendency (p=0.07) toward increased interleukin-6 concentrations was observed.

Discussion: We found adverse acute effects with regard to symptoms, signs, PEF, and inflammatory markers in healthy volunteers exposed 3 hours to diesel (300 µg/m³ and NO₂ 1.4 ppm). TLV(8hours) of NO₂ is 1 ppm. These effects were first seen after 75 min of exposure. Preliminary results show correlations between heart rate variability and sound pressure.

NR 2011;45(5)

55th Nordic Work Environment Meeting
(Nordiska Arbetsmiljömötet)

The Work Environment – Impact of Technological, Social and Climate Change

*Editors: Maria Albin, Johanna Alkan-Olsson, Mats Bohgard,
Kristina Jakobsson, Björn Karlson, Peter Lundqvist,
Mikael Ottosson, Fredrik Rassner, Måns Svensson, and
Håkan Tinnerberg.*

VETENSKAPLIG SKRIFTSERIE

ARBETE OCH HÄLSA

NR 2011;45(5)

55th Nordic Work Environment Meeting
(Nordiska Arbetsmiljömötet)

The Work Environment – Impact of Technological, Social and Climate Change

*Editors: Maria Albin, Johanna Alkan-Olsson, Mats Bohgard,
Kristina Jakobsson, Björn Karlson, Peter Lundqvist,
Mikael Ottosson, Fredrik Rassner, Måns Svensson, and
Håkan Tinnerberg.*

ARBETE OCH HÄLSA

ISBN 978-91-85971-32-9

VETENSKAPLIG SKRIFTSERIE

ISSN 0346-7821