

# LUND UNIVERSITY

# FLT3-ITDs Instruct a Myeloid Differentiation and Transformation Bias in Lymphomyeloid Multipotent Progenitors

Mead, Adam J.; Kharazi, Shabnam; Atkinson, Deborah; Macaulay, Iain; Pecquet, Christian; Loughran, Stephen; Lutteropp, Michael; Woll, Petter; Chowdhury, Onima; Luc, Sidinh; Buza-Vidas, Natalija; Ferry, Helen; Clark, Sally-Ann; Goardon, Nicolas; Vyas, Paresh; Constantinescu, Stefan N.; Sitnicka Quinn, Ewa; Nerlov, Claus; Jacobsen, Sten Eirik W. Published in: Cell Reports

DOI: 10.1016/j.celrep.2013.04.031

2013

Link to publication

Citation for published version (APA):

Mead, A. J., Kharazi, S., Atkinson, D., Macaulay, I., Pecquet, C., Loughran, S., Lutteropp, M., Woll, P., Chowdhury, O., Luc, S., Buza-Vidas, N., Ferry, H., Clark, S.-A., Goardon, N., Vyas, P., Constantinescu, S. N., Sitnicka Quinn, E., Nerlov, C., & Jacobsen, S. E. W. (2013). FLT3-ITDs Instruct a Myeloid Differentiation and Transformation Bias in Lymphomyeloid Multipotent Progenitors. *Cell Reports*, *3*(6), 1766-1776. https://doi.org/10.1016/j.celrep.2013.04.031

Total number of authors: 19

#### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

You may not further distribute the material or use it for any profit-making activity or commercial gain
You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

**PO Box 117** 221 00 Lund +46 46-222 00 00





# FLT3-ITDs Instruct a Myeloid Differentiation and Transformation Bias in Lymphomyeloid Multipotent Progenitors

Adam J. Mead,<sup>1,\*</sup> Shabnam Kharazi,<sup>3</sup> Deborah Atkinson,<sup>1</sup> Iain Macaulay,<sup>1,2</sup> Christian Pecquet,<sup>4,5</sup> Stephen Loughran,<sup>1</sup> Michael Lutteropp,<sup>1,2</sup> Petter Woll,<sup>1</sup> Onima Chowdhury,<sup>1</sup> Sidinh Luc,<sup>1</sup> Natalija Buza-Vidas,<sup>1</sup> Helen Ferry,<sup>1</sup> Sally-Ann Clark,<sup>1</sup> Nicolas Goardon,<sup>2</sup> Paresh Vyas,<sup>2</sup> Stefan N. Constantinescu,<sup>4,5</sup> Ewa Sitnicka,<sup>3</sup> Claus Nerlov,<sup>2,6</sup> and Sten Eirik W. Jacobsen<sup>1,2,\*</sup>

<sup>1</sup>Haematopoietic Stem Cell Biology Laboratory

<sup>2</sup>MRC Molecular Haematology Unit

Weatherall Institute of Molecular Medicine, University of Oxford, Oxford OX3 9DS, UK

<sup>3</sup>Hematopoietic Stem Cell Laboratory, Lund Stem Cell Center, Lund University, Lund 22184, Sweden

<sup>4</sup>Ludwig Institute for Cancer Research, Brussels B1200, Belgium

<sup>5</sup>de Duve Institute, Université Catholique de Louvain, Brussels B1200, Belgium

<sup>6</sup>Institute for Stem Cell Research, MRC Centre for Regenerative Medicine, University of Edinburgh, King's Buildings, West Mains Road, Edinburgh EH93JQ, UK

\*Correspondence: adam.mead@imm.ox.ac.uk (A.J.M.), sten.jacobsen@imm.ox.ac.uk (S.E. W.J.)

http://dx.doi.org/10.1016/j.celrep.2013.04.031

### SUMMARY

Whether signals mediated via growth factor receptors (GFRs) might influence lineage fate in multipotent progenitors (MPPs) is unclear. We explored this issue in a mouse knockin model of gain-offunction Flt3-ITD mutation because FLT3-ITDs are paradoxically restricted to acute myeloid leukemia even though Flt3 primarily promotes lymphoid development during normal hematopoiesis. When expressed in MPPs, Flt3-ITD collaborated with Runx1 mutation to induce high-penetrance aggressive leukemias that were exclusively of the myeloid phenotype. Flt3-ITDs preferentially expanded MPPs with reduced lymphoid and increased myeloid transcriptional priming while compromising early B and T lymphopoiesis. Flt3-ITD-induced myeloid lineage bias involved upregulation of the transcription factor Pu.1, which is a direct target gene of Stat3, an aberrantly activated target of FIt3-ITDs, further establishing how lineage bias can be inflicted on MPPs through aberrant GFR signaling. Collectively, these findings provide new insights into how oncogenic mutations might subvert the normal process of lineage commitment and dictate the phenotype of resulting malignancies.

#### INTRODUCTION

Whether signals mediated via growth factor receptors (GFRs) might influence lineage fate in normal multipotent progenitors (MPPs) and stem cells remains unclear and disputed. New in-

sights into this process have recently been gained through studies in vitro (Rieger et al., 2009); however, whether GFR signaling instructs lineage specification in vivo remains a key unresolved issue (Enver and Jacobsen, 2009). Studying the impact of gain-of-function mutations is an alternative approach to determine whether and how GFRs might instruct lineage fate in vivo. Indeed, activating mutations of GFRs and downstream signaling pathways are common events in cancer, particularly in hematopoietic malignancies such as acute myeloid leukemia (AML), and often show striking associations with distinct clinical and celllineage phenotypes (Croce, 2008). In some cases, this might simply reflect the fact that such mutations primarily target a specific cell lineage. However, more intriguingly, if a mutation targets a primitive multipotent cell, it might instruct lineage-fate decisions. As recent investigations of human AML have suggested that the propagating cell might frequently represent the counterpart of normal MPPs (Goardon et al., 2011), it is possible but still unclear to what degree GFR signaling mutations might also influence lineage specification in multipotent cells.

A good example of aberrant GFR signaling associated with specific leukemia phenotypes relates to the FMS-like tyrosine kinase 3 receptor (FLT3), which is expressed in the majority of cases of AML and acute lymphoblastic leukemia (ALL) (Stirewalt and Radich, 2003). Constitutively activating internal tandem duplications (ITDs) within the juxtamembrane domain of FLT3 occur in ~25% of cases of AML, conferring an adverse prognosis (Stirewalt and Radich, 2003). However, despite the high frequency of FLT3 expression in ALL and the key role of Flt3 in early lymphoid development (Sitnicka et al., 2002), including high-level Flt3 expression in lymphoid-primed MPPs (LMPPs) with combined lymphoid and myeloid potential (Adolfsson et al., 2005), FLT3-ITDs are rare in cases of ALL, occurring in <1% in larger series (Leow et al., 2011).

As is the case for most leukemic mutations, it is unclear how often FLT3-ITDs are a true initiating event in myeloid





malignancies. Indeed, a number of lines of evidence support the notion that FLT3-ITDs are often acquired secondarily to an initiating clonogenic event (Gale et al., 2008; Jan et al., 2012), although when involved in chronic myelomonocytic leukemia (CMML), they may indeed be the initiating mutation (Lee et al., 2007). Nevertheless, in line with the normal expression pattern of Flt3 (Adolfsson et al., 2005), results from studies in patients are compatible with the notion that FLT3-ITDs occur within the human MPP compartment (Levis et al., 2005) and are an essential driver mutation within the founding leukemic clone (Ding et al., 2012; Smith et al., 2012). Thus, even though FLT3-ITDs might occur secondarily to an initiating clonogenic event in many cases, these mutations occur in multipotent cells (Levis et al., 2005) and FLT3-ITD GFR signaling appears to be an essential requirement for leukemia propagation (Smith et al., 2012). It is possible, therefore, that FLT3-ITDs may act to dictate the lineage fate and phenotype of the resulting leukemia. Compatible with a role of Flt3-ITD signaling in lineage determination, two different knockin mouse models of Flt3-ITD have been reported to develop a myeloproliferative phenotype exclusively (Lee et al., 2007; Li et al., 2008). However, although it was recently suggested that Flt3-ITDs deplete hematopoietic stem cells (HSCs) (Chu et al., 2012), the key progenitor population that propagates FLT3-ITD-induced myeloid disease, as well as the cellular and molecular bases of their myeloid lineage bias, remains unclear.

Using a mouse knockin model of the *Flt3-ITD* mutation, we investigated the cellular and molecular mechanisms by which constitutive GFR signaling might subvert lineage specification in MPPs and alter the cell fate of early lymphoid progenitors, in order to explain the myeloid bias of the resulting leukemias.

#### RESULTS

# Flt3-ITD Collaborates with Runx1 Mutation to Induce Aggressive AML

To definitively determine whether physiologically expressed FLT3-ITD impacts the establishment of myeloid versus lymphoid leukemia development, we crossed Flt3<sup>ITD/ITD</sup> with Runx1<sup>fl/fl</sup> (Growney et al., 2005) and Mx1-Cre mice to induce Runx1 deletion in MPPs. Importantly, Runx1 loss-of-function mutation is associated with both lymphoid and myeloid leukemia (Grossmann et al., 2011; Schnittger et al., 2011). Unexpectedly, even without poly I:C induction, Flt3<sup>/TD/ITD</sup>Runx1<sup>fl/fl</sup>Mx1-Cre<sup>+</sup> mice developed a high-penetrance, short-latency acute leukemia (Figure 1A) characterized by marked leukocytosis, anemia, and thrombocytopenia (Figures S1A-S1C) and hepatosplenomegaly. Peripheral blood (PB) and bone marrow (BM) morphology resembled AML in all diseased mice (Figures 1B-1D), and myeloid lineage was confirmed by flow cytometry (Figures 1E and S1D). Development of ALL was never observed. Leukemias showed deletion of the Runx1 DNA-binding domain (Figure S1E). The uniform myeloid-lineage leukemias in Flt3'TD/ITD Runx1<sup>fl/fl</sup> Mx1-Cre<sup>+</sup> mice demonstrate a key role for Flt3-ITD signaling in introducing a myeloid-lineage bias during leukemogenesis.

### Flt3-ITDs Expand Myeloid-Biased LMPPs

We next investigated the cellular and molecular bases for FLT3-ITD-induced myeloid bias. As shown previously (Lee

et al., 2007), the multipotent Lin-Sca1+c-Kit+ (LSK) compartment was expanded in *Flt3<sup>ITD/ITD</sup>* mice (Figure 2A). We applied CD150 and CD48 to determine the nature of the expanded cells within the LSK compartment (Kiel et al., 2005), as Flt3-ITD is not detectable at the cell surface (Figure S2A). Notably, the expansion of LSKs was wholly attributable to a marked expansion of LSKCD150<sup>-</sup>48<sup>+</sup> MPPs in 8- to 10-week-old Flt3<sup>/TD/ITD</sup> mice (Figures 2B and 2C), in line with the recent observation that Flt3-ITD may suppress the HSC compartment (Chu et al., 2012). Heterozygous (Flt3/TD/+) mice had an intermediate phenotype between  $Flt3^{+/+}$  and  $Flt3^{ITD/ITD}$  mice (Figures S2B and S2C). High-level Flt3 expression in the normal LSK compartment defines LMPPs that lack self-renewal and megakaryocytic (Mk) and erythroid (E) potential but sustain lymphomyeloid capability (Adolfsson et al., 2005), a progenitor that is also implicated in human AML (Goardon et al., 2011). Because Flt3<sup>high</sup> LMPPs reside almost exclusively in the LSKCD150<sup>-48<sup>+</sup></sup> compartment (Figure S2D), we explored whether the expanded LSKCD150<sup>-48<sup>+</sup></sup> cells in *Flt3-ITD* mice had LMPP-like characteristics. As expected for LMPPs (Mansson et al., 2007), HSC- and MkE-affiliated gene expression was downregulated in Flt3<sup>ITD/ITD</sup> LSKCD150<sup>-48<sup>+</sup></sup> MPPs (Figures 2D, 2E, and S2E), and in line with the molecular data they possessed little or no Mk potential in vitro (Figure 2F). In contrast, myeloid-affiliated gene expression was upregulated in Flt3<sup>ITD/ITD</sup> LSKCD150-48<sup>+</sup> cells (Figure 2G), paralleled by high granulocyte-macrophage (GM) potential in vitro (Figure 2H). Early lymphoid transcriptional programs were downregulated in both  $Flt3^{ITD/ITD}$  and  $Flt3^{ITD/+}$ LSKCD150<sup>-48<sup>+</sup></sup> MPPs (Figures 2I and 2J), paralleled by severely reduced B cell potential in vitro (Figure 2K). Importantly, the lymphoid transcriptional program was already suppressed in Flt3<sup>ITD/ITD</sup> LSKCD150<sup>-</sup>48<sup>+</sup> MPPs in embryonic day 15 (E15) fetal liver (FL) (Figure S2F), at which time the phenotype and number of LSKCD150<sup>-48<sup>+</sup></sup> MPPs were unaffected (Figure S2G). In keeping with this suppression of early lymphoid programs in fetal MPPs, the number of B220<sup>+</sup>CD19<sup>+</sup> B cells was suppressed in Flt3-ITD E15 FL (Figure S2H), preceding the emergence of a myeloproliferative phenotype at this early stage of ontogeny (Figure S2I). This supports the notion that lymphoid suppression and myeloid bias in MPPs occurs as a cell-intrinsic and direct consequence of Flt3-ITD signaling.

Because FLT3-ITDs in humans often occur secondarily to an initiating clonogenic event (Jan et al., 2012), it is possible that in such cases it is the other genetic events and not the FLT3-ITD that influence the leukemic phenotype. In order to address this issue, we used Vav-Cre mediated recombination to examine the impact of Runx1 loss of function on lineage priming in MPPs in the absence of Flt3-ITD. The data obtained demonstrate a significant upregulation of *II7r* and *Rag1* expression in MPPs with no significant impact on slgH expression (Figure S2J). This finding is in keeping with the high incidence of lymphoid malignancies in mouse models of Runx1 mutation (Jacob et al., 2010; Kundu et al., 2005; Putz et al., 2006) and contrasts markedly with the suppression of lymphoid programs caused by Flt3-ITDs in MPPs. This supports the notion that it is GFR signaling through Flt3-ITD that instructs the uniform myeloid phenotype of leukemias resulting from the collaboration between Runx1 deletion and FLT3-ITD in LMPPs.





Figure 1. Mutation of *Runx1* in *Flt3*<sup>/TD/ITD</sup> Mice Results in High-Penetrance Aggressive Myeloid Leukemia

(A) Leukemia-free survival curves of Flt3<sup>ITD/ITD</sup>Runx1<sup>fl/fl</sup> mice stratified according to Mx1Cre genotype (n = 50–68 mice of each genotype).

(B and C) Low-power (20×; B) and high-power (100×; C) morphology of typical leukemic cells in *Flt3<sup>TD/ITD</sup>Runx1<sup>#/#</sup>Mx1Cre*<sup>+</sup> mice. (D) Differential morphology results from BM of WT (+/+), *Flt3<sup>TD/ITD</sup>Runx1<sup>#/#</sup>Mx1Cre*<sup>-</sup>, and leukemic *Flt3<sup>TD/ITD</sup>Runx1<sup>#/#</sup>Mx1Cre*<sup>+</sup> mice; percentage of total BM cells. Mean (SEM) values for 4-11 mice of each genotype.

(E) Fluorescence-activated cell sorting (FACS) analysis of PB and BM from an Flt3<sup>/TD/ITD</sup>Runx1<sup>fl/fl</sup>Mx1Cre<sup>+</sup> mouse with myeloid leukemia (representative of 19 further analyzed cases).

Error bars represent SEM. See also Figure S1.





## Figure 2. Flt3-ITDs Expand Myeloid-Biased MPPs

(A) Expansion of LSK cells in 8- to 10-week-old *Flt3<sup>/TD/ITD</sup>* mice (+/+: n = 27; ITD/ITD: n = 23).

(B) Expression of CD48 and CD150 on BM LSK cells from *Flt3*<sup>+/+</sup> and *Flt3*<sup>ITD/ITD</sup> mice.

 (C) Expansion of LSKCD48<sup>+</sup>CD150<sup>-</sup> MPPs in 8- to 10-week-old *Flt3<sup>(TD/ITD</sup>* mice (+/+: n = 27; ITD/ITD: n = 23).
(D) GSEA demonstrating downregulation of E-affiliated genes in *Flt3<sup>(TD/ITD</sup>* versus *Flt3<sup>+/+</sup>LSKCD48<sup>+</sup>150<sup>-</sup>* MPPs (n = 4 of each genotype). FDR, false discovery rate; NES, normalized enrichment score.

(legend continued on next page)



# FIt3-ITD Suppresses Early T- and B-Lymphoid Progenitors

Because Flt3-ITD compromised lymphoid-transcriptional priming in LMPPs, which have been implicated as critical thymusseeding progenitors (Luc et al., 2012), we next investigated whether T lymphopoiesis might be suppressed in Flt3-ITD mice. Thymic cellularity was found to be progressively reduced, and the earliest thymic progenitors (double-negative 1 Kit+ [DN1Kit<sup>+</sup>]) were almost completely lost in *Flt3<sup>ITD/ITD</sup>* mice (Figures 3A-3C). Also, subsequent stages of DN2Kit<sup>+</sup> and DN3 thymocyte progenitors were severely reduced (Figures S3A and S3B). Of relevance, gene and protein expression for the chemokine receptor Ccr9, which is critical for migration of thymus-seeding progenitors from the BM to the thymus (Schwarz et al., 2007), was suppressed in Flt3<sup>ITD/ITD</sup> LSKCD150<sup>-48+</sup> MPPs (Figures 3D, S3C, and S3D). Moreover, DN1 cells showed large reductions in expression of Notch1 and its targets (Hes1 and Hes5; Figure S3E), which are critical for early T cell development. Although expression of some early lymphoid genes (II2ra, II7r, and Gata3) was maintained, myeloid genes (Cebpa, Sfpi1, Csf1r, Csf2r, and Csf3r) were highly upregulated in DN1 thymic progenitors (Figure S3E). In keeping with this myeloid bias, Flt3<sup>ITD/ITD</sup> DN1 progenitors showed low T cell potential in vitro (Figure 3E).

Previous studies demonstrated a reduction of mature B cells in Flt3-ITD mice (Lee et al., 2007; Li et al., 2008). We investigated whether this might reflect Flt3-ITD-induced perturbation at the earliest B cell commitment stages, and in support of this found a notable 11-fold expansion of Lin<sup>-</sup>CD19<sup>-</sup>CD24<sup>-</sup> AA4.1<sup>+</sup>CD43<sup>mid</sup>B220<sup>+</sup> pre-pro-B cells in *Flt3<sup>ITD/ITD</sup>* mice (Figures 3F and 3G), whereas pro-B cells, the next stage in B cell development, were severely reduced (Figures 3H and S3F). Because pre-pro-B cells are known to sustain low-level myeloid potential (Rumfelt et al., 2006), we next investigated lymphoid and myeloid gene expression in Flt3<sup>ITD/ITD</sup> pre-pro-B cells. Key early lymphoid genes (Cd79a, II7r, slgh, Ebf1, Pax5, Rag1, and Rag2) were all suppressed (Figure 3I), whereas myeloid-affiliated genes (Cebpa, Cebpb, Sfpi1, Csf1r, and Csf2r) were upregulated in Flt3/ITD/ITD pre-pro-B cells (Figure 3J). In agreement with this, Flt3/ITD/ITD pre-pro-B cells had severely reduced B cell potential (Figure 3K). These findings demonstrate that FIt3-ITDs markedly impair the earliest stage of both T and B lymphopoiesis and upregulate myeloid gene expression in the earliest B and T lymphoid progenitors, in agreement with the notion that myeloid bias initiates in multipotent LMPPs.

# Flt3-ITD-Induced Myeloid Bias Is Dependent on Upregulation of Pu1

To further explore the mechanistic basis of Flt3-ITD-induced myeloid bias of early lymphoid progenitors, we examined Pu1 protein expression using a Pu1-YFP reporter line. Whereas in wild-type (WT) mice Lin<sup>-</sup> cells showed a bimodal distribution between Pu1<sup>low/-</sup> and Pu1<sup>high</sup> cells, almost all Lin<sup>-</sup> cells in Flt3<sup>ITD/ITD</sup> mice were Pu1<sup>high</sup> (Figure 4A). Flt3<sup>ITD/ITD</sup> mice showed a modest increase in Pu1 expression in LSKCD150<sup>-48+</sup> MPPs (Figure 4B) and markedly enhanced levels of Pu1-YFP in prepro-B cells (Figure 4C). In keeping with the observed increased expression of Pu1, gene set enrichment analysis (GSEA) demonstrated upregulation of Pu1 target genes (Steidl et al., 2006) in *Flt3-ITD* LSKCD150<sup>-</sup>48<sup>+</sup> MPPs (Figure 4D). Furthermore, GSEA also demonstrated a similar upregulation of Pu1 target genes in FLT3-ITD mutated "LMPP-like" leukemia stem cells in human AML (Goardon et al., 2011) in comparison with FLT3-WT counterparts (Figure S4A).

Because transition from pre-pro-B cells to subsequent stages of B cell development is associated with downregulation of Pu1 expression (Figure S4B), we attempted to rescue the suppressed B cell phenotype in Flt3-ITD mice through generation of *Flt3<sup>ITD/ITD</sup>* Pu1<sup>+/-</sup> mice. Notably, whereas Pu1 haploinsufficiency did not increase the number of B cells or their progenitors on a WT background, it resulted in a 10.3-fold increase in B cells in *Flt3<sup>ITD/ITD</sup>* mice (Figure 4E). Furthermore, Pu1 haploinsufficiency led to a 2fold reduction in the expanded pre-pro-B cell population in Flt3-ITD mice (Figure 4F) together with a 13.5-fold rescue of the suppression of pro-B cells (Figure 4G). Strikingly, in the presence of Pu1 haploinsufficiency, the expansion of MPPs was also restored to WT levels (Figure 4H) together with a significant amelioration of the myeloproliferative phenotype in *Flt3<sup>ITD/ITD</sup>* mice (Figure 4I).

Stat3 is aberrantly activated by Flt3-ITD signaling (Schmidt-Arras et al., 2009) and Pu1 is a direct target gene of Stat3 (Hegde et al., 2009). In line with this, *Flt3<sup>ITD/ITD</sup>* BM showed aberrant activation of STAT3 (Figure S4C), and MPPs and pre-pro-B cells showed significantly increased expression of Stat3 target genes (Figures 4J–4L). Collectively, these data suggest that Pu1 upregulation is a key mediator of the Flt3-ITD-induced myeloid bias and progenitor phenotype.

### DISCUSSION

In the studies presented here, we explored whether and how GFR signaling might influence lineage specification in vivo. To

Error bars represent SEM. See also Figure S2.

<sup>(</sup>E) Reduced MkE-affiliated gene expression in LSKCD48<sup>+</sup>150<sup>-</sup> cells from *Flt3<sup>+/+</sup>*, *Flt3<sup>ITD/+</sup>*, and *Flt3<sup>ITD/+D</sup>* mice relative to *Flt3<sup>+/+</sup>* LSK FLT3<sup>-</sup> cells; p values represent a comparison of *Flt3<sup>+/+</sup>* LSK FLT3<sup>-</sup> cells versus LSKCD48<sup>+</sup>150<sup>-</sup> *Flt3<sup>ITD/ITD</sup>* cells by unpaired t test (n = 6–8 replicates from 2–3 mice per genotype). Differences among *Flt3<sup>+/+</sup>*, *Flt3<sup>ITD/+</sup>*, and *Flt3<sup>ITD/ITD</sup>* MPPs were not significant.

<sup>(</sup>F) Single-cell Mk potential of Flt3<sup>+/+</sup>and Flt3<sup>+/+</sup>and Flt3<sup>+/+</sup>and Flt3<sup>+/+</sup>LSKCD48<sup>+</sup>150<sup>-</sup> MPPs in comparison with Flt3<sup>+/+</sup>LSKCD48<sup>-</sup>150<sup>+</sup> HSCs (n = 3 independent experiments).

<sup>(</sup>G) GSEA demonstrating upregulation of myeloid affiliated genes in *Flt3<sup>ITD/ITD</sup>* versus *Flt3<sup>+/+</sup>*LSKCD48<sup>+</sup>150<sup>-</sup> MPPs (n = 4 of each genotype).

<sup>(</sup>H) Single-cell GM potential of  $Flt3^{+/+}$  and  $Flt3^{1/D/1TD}$  LSKCD48<sup>+</sup>150<sup>-</sup> MPPs. Open bars show the frequency of clones formed, and closed bars show the frequency of highly proliferative clones (covering >50% of the well; n = 3 independent experiments).

<sup>(</sup>I) GSEA demonstrating downregulation of lymphoid transcriptional program in *Flt3<sup>T/D/ITD</sup>* versus *Flt3<sup>+/+</sup>* LSKCD48<sup>+</sup>150<sup>-</sup> MPPs (n = 4 of each genotype).

<sup>(</sup>J) Expression of lymphoid-affiliated genes in LSKCD48<sup>+150<sup>-</sup></sup> MPPs from *Flt3<sup>+/+</sup>*, *Flt3<sup>TD/+</sup>*, and *Flt3<sup>TD/TD</sup>* mice (n = 6–8 replicates from 2–3 mice per genotype); p values indicate comparisons of *Flt3<sup>+/+</sup>* versus *Flt3<sup>TD/+</sup>* MPPs, and *Flt3<sup>+/+</sup>* versus *Flt3<sup>TD/TD</sup>* MPPs by unpaired t test (p < 0.001 for all analyses).

<sup>(</sup>K) B cell potential (B220<sup>+</sup>CD19<sup>+</sup>) of *Flt3<sup>+/+</sup>* versus *Flt3<sup>/TD//TD</sup>* BM LSKCD48<sup>+</sup>150<sup>-</sup> MPPs (10 cells/well). Data are mean (SEM) values of 2–3 experiments, with 9–12 replicate wells in each experiment.





(legend on next page)

that end, we used a knockin model of *Flt3-ITD* to investigate whether oncogenic mutations that result in constitutive GFR signaling influence the lineage fate of MPPs in vivo.

In order to first establish the physiological relevance of Flt3-ITD as an AML-inducing mutation, we developed a mouse model in which Flt3-ITD was coexpressed with an inducible deletion of the DNA-binding domain of Runx1 (Growney et al., 2005) in MPPs. Importantly, although Runx1 loss-of-function mutations are found in patients with both ALL and AML, only the latter is often found in association with FLT3-ITDs (Grossmann et al., 2011; Schnittger et al., 2011). Deletion of the DNA-binding domain of Runx1 during adult murine hematopoiesis results in mild myeloproliferation (Growney et al., 2005) or lymphoblastic leukemias/lymphomas (Jacob et al., 2010; Kundu et al., 2005; Putz et al., 2006). Here, we demonstrate that the combination of Runx1 mutation with Flt3-ITDs results in aggressive leukemia with 100% penetrance and a uniform myeloid phenotype. In addition to demonstrating the functional relevance for FLT3-ITD-induced AML, this AML model has a number of important features. First, in contrast to many other mutation collaboration models, both mutations are targeted to the physiologically relevant endogenous genetic loci. Second, in isolation each mutation results in only a modest phenotype, but in combination a minor clone of deleted cells rapidly expands and becomes clonally dominant, paralleling somatic mutations during leukemogenesis. Third, the evolution of leukemia is rapid, supporting the notion that it occurs without the requirement for additional genetic events. Finally, the model recapitulates a particularly poor prognostic form of AML with a major unmet need for novel therapeutic approaches and thus provides a powerful model for future studies of the cellular and molecular mechanistic bases for collaboration of these mutations in AML.

The most primitive progenitor population that was expanded by Flt3-ITDs consisted of LMPP-like cells, demonstrating upregulation of the myeloid program, whereas the transcriptional expression of lymphoid-affiliated genes was markedly reduced in both heterozygous and homozygous Flt3-ITD mice. This myeloid transcriptional bias and suppression of lymphoid transcripts was also present during early B and T cell development and, importantly, was already present in Flt3-ITD FL MPPs, preceding the expansion of MPPs and development of a myeloproliferative phenotype, supporting the notion that it is a direct consequence of Flt3-ITD signaling in LMPPs. The severe suppression of the earliest thymic progenitors in *Flt3-ITD* mice and concomitant lack of Ccr9 upregulation in LMPPs, which have been implicated as key thymus-seeding progenitors (Luc et al., 2012), further suggests that thymic seeding might also be impaired by Flt3-ITDs. These findings demonstrate that the myeloid propensity of FLT3-ITDs results from FLT3-ITD introducing a myeloid bias in multipotent lymphomyeloid progenitors as well as in the earliest B and T lymphoid progenitors.

Pu1 is a dosage-sensitive regulator of myeloid-lymphoid cell-fate decisions that promotes myeloid differentiation when overexpressed (DeKoter and Singh, 2000). Using a Pu1-YFP reporter, we demonstrated that Flt3-ITDs upregulate Pu1 expression in MPPs and the earliest B lymphoid progenitors, paralleled by increased expression of Pu1 target genes in MPPs. Furthermore, we confirmed the relevance of these findings for human AML by demonstrating upregulation of Pu1 target genes in FLT3-ITD mutated LMPP-like human AML stem cells (Goardon et al., 2011). In agreement with a key mechanistic role for Pu1 overexpression, Flt3-ITD-induced suppression of B cell development, as well as the Flt3-ITD-associated progenitor phenotype, was partially rescued by Pu1 haploinsufficiency.

Constitutive Flt3-ITD signaling is distinct from WT Flt3 signaling due to abnormal anchoring of FLT3-ITD in the endoplasmic reticulum, with markedly reduced surface Flt3 expression resulting in aberrant Stat3 activation, as confirmed in our study (Schmidt-Arras et al., 2009). Because Pu1 is a direct target gene of Stat3 (Hegde et al., 2009) and Stat3 target genes were upregulated in *Flt3-ITD* LSKCD150<sup>-48+</sup> MPPs and pre-pro-B cells, this provides a putative direct link between aberrant constitutive Flt3-ITD signaling and Pu1 overexpression as a cause of aberrant myeloid bias.

Several lines of evidence indicate that FLT3-ITDs frequently arise early in the transformation process in MPPs (Levis et al., 2005) and as a driver mutation in the founding leukemic clone (Ding et al., 2012; Smith et al., 2012), and thus may influence the lineage of resulting leukemias. Together with our findings,

(C) Progressive reduction of DN1Kit<sup>+</sup> cells in *Flt3<sup>/TD/ITD</sup>* thymus.

Figure 3. Early T and B Cell Progenitors Are Suppressed and Myeloid Biased in *Flt3<sup>(TD/ITD</sup>* Mice

<sup>(</sup>A) Thymic cellularity of Flt3<sup>+/+</sup> and Flt3<sup>ITD/ITD</sup> mice at 2–3 and 8–10 weeks (n = 7–10 mice of each genotype at each age).

<sup>(</sup>B) Representative DN staging of Flt3<sup>+/+</sup> and Flt3<sup>/TD/ITD</sup> thymocytes from 2-week-old mice; percentages are relative to total thymocytes.

<sup>(</sup>D) Expression (SEM) of *Ccr9* in LSKCD48<sup>+</sup>150<sup>-</sup> BM cells from 8- to 10-week-old *Flt3<sup>+/+</sup>* and *Flt3<sup>ITD/ITD</sup>* mice (n = 3–4 mice per genotype, 2–3 separately sorted replicates per mouse).

<sup>(</sup>E) T cell potential on OP9DL1 cells of DN1 cells (10/cells per well) from 2- to 3-week-old *Flt3<sup>+/+</sup>* and *Flt3<sup>ITD/ITD</sup>* mice (n = 2 mice per genotype, 12 replicate wells per mouse). ND, not detected.

<sup>(</sup>F) Representative staging of early B cell progenitors in *Flt3<sup>+/+</sup>* and *Flt3<sup>TD/TD</sup>* 2- to 3-week-old mice. pre-pro-B cells are identified as Lin<sup>-</sup>B220<sup>+</sup>CD19<sup>-</sup>CD24<sup>-</sup>AA4.1<sup>+</sup>CD43<sup>mid</sup>; percentage of total BM cells.

<sup>(</sup>G) Increased numbers of pre-pro-B cells in 2- to 3-week-old Flt3/TD/ITD mice (mean [SEM] values for 2- to 3-week-old mice, n = 8-10 of each genotype).

<sup>(</sup>H) Reduced numbers of pro-B cells (Lin<sup>-</sup>B220<sup>+</sup>CD43<sup>+</sup>CD19<sup>+</sup>CD24<sup>int</sup>AA4.1<sup>+</sup>) in *Flt3<sup>ITD/ITD</sup>* mice.

<sup>(</sup>I) Mean (SEM) expression of lymphoid-affiliated genes in pre-pro-B cells from 2- to 3-week-old mice (n = 2 mice per genotype, 3 separately sorted replicates per mouse).

<sup>(</sup>J) Mean (SEM) expression of myeloid-affiliated genes in pre-pro-B cells from *Flt3*<sup>/TD/ITD</sup> mice.

<sup>(</sup>K) In vitro B cell potential (B220<sup>+</sup>CD19<sup>+</sup> cells) of pre-pro-B cells from 2- to 3-week-old mice (n = 10 cells/well, 2 mice per genotype, 10 replicate wells per mouse). Error bars represent SEM. See also Figure S3.





(legend on next page)



this is consistent with a model in which LMPPs, which normally express high levels of FLT3 (Adolfsson et al., 2005), acquire FLT3-ITD mutations that confer both a strong clonal advantage and a marked myeloid bias. This results in myeloid expansion and suppression of early lymphoid development, strongly supporting a fundamental role for FLT3-ITDs in promoting myeloid lineage leukemia development at the MPP level. Our study also further highlights LMPPs as a key target population in AML, as also supported by recent findings in human AML (Goardon et al., 2011). Transformation to an aggressive leukemia exclusively of a myeloid phenotype by introduction of Runx1 mutation demonstrates the functional relevance of this FLT3-ITD-induced myeloid bias and clonal dominance, providing insights into the process by which oncogenic mutations might determine the lineage fate of the resulting leukemias at the precommitment stage. Our findings are also of considerable relevance for normal hematopoiesis, as it remains disputed whether key cytokine receptor signaling pathways mediate critical in vivo functions in blood lineage development through purely permissive rather than instructive actions (Enver and Jacobsen, 2009). Thus, although Flt3-ITD elicits aberrant signaling, our findings clearly provide support for the notion that cytokine receptors are also capable of eliciting lineage-instructive signaling in MPPs in vivo.

#### **EXPERIMENTAL PROCEDURES**

#### Animals

All animals used were bred and maintained in accordance with regulations of the UK Home Office. Details regarding the mouse lines are provided in the Extended Experimental Procedures.

#### **Patient Samples**

BM samples from AML patients were obtained with informed consent and the approval of Oxford Ethics Committee B (protocols 06/Q1606/110 and 05/ MRE07/74).

#### Fluorescence-Activated Cell Sorting

Details of the antibody staining panels and protocols are provided in the Extended Experimental Procedures. Antigens and antibodies used for identification of specific cell populations are shown in Tables S1 and S2.

#### Western Blotting

Cells were lysed in 1% NP40 buffer and analyzed by western blot as previously described (Pecquet et al., 2010) with antibodies against pStat3 (Cell Signaling) and beta-actin (Sigma).

#### In Vitro Evaluation of Lineage Potentials

Evaluation of megakaryocytic, GM, and lymphoid potentials was carried out as previously described (Månsson et al., 2007) and detailed in the Extended Experimental Procedures.

#### Gene Expression by Dynamic Arrays

Multiplex quantitative PCR was performed as previously described (Kharazi et al., 2011) and detailed in the Extended Experimental Procedures. Assays used for dynamic arrays are shown in Table S3.

#### **Microarray Analysis**

Analysis by Affymetrix Mouse Genome 430 2.0 Arrays was carried out at the Stanford Protein and Nucleic Acid Facility as described previously (Månsson et al., 2007) and in the Extended Experimental Procedures.

#### **ACCESSION NUMBERS**

The GEO accession number for the microarray data reported in this paper is GSE35805.

#### SUPPLEMENTAL INFORMATION

Supplemental Information includes Extended Experimental Procedures, four figures, and three tables and can be found with this article online at http://dx.doi.org/10.1016/j.celrep.2013.04.031.

#### LICENSING INFORMATION

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-No Derivative Works License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

#### ACKNOWLEDGMENTS

We thank Gary Gilliland for kindly providing Flt3-ITD knockin mice. A.M. and D.A. were funded by a Leukaemia and Lymphoma Research Senior Bennett Fellowship, S.K. was funded by the Pasteur Institute of Iran, and S.E.W.J. was funded through a Strategic Appointment and Programme grant from the

### Figure 4. Involvement of PU1 Upregulation in the Myeloid Bias of Flt3<sup>/TD/ITD</sup> Mice

(A–C) Representative FACS analysis of PU1-YFP expression in lineage-negative (A), LSKCD48<sup>+</sup>150<sup>-</sup> MPPs (B), and Lin<sup>-</sup>CD19<sup>-</sup>CD24<sup>-</sup>AA4.1<sup>+</sup>CD43<sup>mid</sup>B220<sup>+</sup> prepro-B (C) cells in BM of 2- to 3-month-old *Flt3<sup>+/+</sup>* and *Flt3<sup>TD/ITD</sup>* mice. Also shown is the mean fold difference in PU1-YFP expression for each population from six mice of each genotype.

(D) GSEA demonstrating upregulation of Pu1 target genes in LSKCD48<sup>+</sup>150<sup>-</sup> MPPs from *Flt3<sup>/TD/ITD</sup>* versus *Flt3<sup>+/+</sup>* mice (n = 4 of each genotype).

(E) Percentage of B220<sup>+</sup>CD19<sup>+</sup> in the BM of 6- to 8-week-old  $Flt3^{+/+}Pu1^{+/+}$ ,  $Flt3^{+/+}Pu1^{+/-}$ ,  $Flt3^{+/+}Pu1^{+/+}$ , and  $Flt3^{+/+}Pu1^{+/-}$  mice (n = 8–11 mice of each genotype; fold difference and p value are for the difference between  $Flt3^{+/}Pu1^{+/+}$  and  $Flt3^{+/}Pu1^{+/-}$  mice).

(F) Percentage of pre-pro-B cells in the BM of 6- to 8-week-old Flt3<sup>+/+</sup>Pu1<sup>+/+</sup>, Flt3<sup>+/+</sup>Pu1<sup>+/-</sup>, Flt3<sup>ITD/ITD</sup>Pu1<sup>+/+</sup> and Flt3<sup>ITD/ITD</sup>Pu1<sup>+/-</sup> mice (n = 3 mice of each genotype).

(G) Percentage of pro-B cells in the BM of 6- to 8-week-old *Flt3<sup>+/+</sup>Pu1<sup>+/+</sup>*, *Flt3<sup>+/+</sup>Pu1<sup>+/-</sup>*, *Flt3<sup>ITD/ITD</sup>Pu1<sup>+/+</sup>* and *Flt3<sup>ITD/ITD</sup>Pu1<sup>+/-</sup>* mice (n = 3 mice of each genotype).

(H) Percentage of LSKCD48<sup>+</sup>150<sup>-</sup> MPPs in the BM of 6- to 8-week-old  $Flt3^{+/+}Pu1^{+/+}$ ,  $Flt3^{+/+}Pu1^{+/-}$ ,  $Flt3^{ITD/ITD}Pu1^{+/+}$  and  $Flt3^{ITD/ITD}Pu1^{+/-}$  mice (n = 7–9 mice of each genotype).

(I) Percentage of Mac1<sup>low</sup>cKit<sup>low</sup> myeloid precursor cells in the BM of 6- to 8-week-old  $Flt3^{+/+}Pu1^{+/+}$ ,  $Flt3^{+/+}Pu1^{+/-}$ ,  $Flt3^{ITD/ITD}Pu1^{+/+}$  and  $Flt3^{ITD/ITD}Pu1^{+/-}$  mice (n = 7–9 mice of each genotype).

(J) GSEA demonstrating upregulation of STAT3 target genes in *Flt3<sup>ITD/ITD</sup>* versus *Flt3<sup>+/+</sup>* LSKCD48<sup>+</sup>150<sup>-</sup> MPPs (n = 4 mice of each genotype).

(K and L) Expression of Stat3 target genes in *Flt3<sup>+/+</sup>* and *Flt3<sup>TD/ITD</sup>* LSKCD48<sup>+150<sup>-</sup></sup> MPPs (K) and pre-pro-B cells (L) (n = 2–4 mice of each genotype, 2–3 replicates per mouse). ND, not detected.

Error bars represent SEM. See also Figure S4.

Medical Research Council (UK) and a Hemato-Linné grant (Swedish Research Council Linnaeus). A.M., P.V., and S.E.J. were supported by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre based at Oxford University Hospitals NHS Trust, and by the NCRN MDSBio sample collection study. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health. A.M. designed, performed, and analyzed experiments and wrote the manuscript. Support to S.N.C. from Fondation contre le cancer, Salus Sanguinis, Action de Recherche Concertée and Interuniversity Attraction Poles Programs and Fonds de la Recherche Scientifique, Belgium and Ludwig Institute for Cancer Research is acknowledged. S.K. performed and analyzed experiments. D.A. provided technical assistance. I.M. and N.G. analyzed microarray data. S. Loughran, M.L., P.W., O.C., S. Luc, N.B.V., H.F., and S.A.C. were involved in FACS analysis/sorting. C.P. performed western blotting. E.S., P.V., S.C., and C.N. provided input on experimental design and analysis. S.E.W.J. conceived and supervised the project, designed and analyzed experiments, and wrote the manuscript.

Received: January 12, 2012 Revised: March 12, 2013 Accepted: April 29, 2013 Published: May 30, 2013

#### REFERENCES

Adolfsson, J., Månsson, R., Buza-Vidas, N., Hultquist, A., Liuba, K., Jensen, C.T., Bryder, D., Yang, L., Borge, O.J., Thoren, L.A., et al. (2005). Identification of Flt3+ lympho-myeloid stem cells lacking erythro-megakaryocytic potential a revised road map for adult blood lineage commitment. Cell *121*, 295–306.

Chu, S.H., Heiser, D., Li, L., Kaplan, I., Collector, M., Huso, D., Sharkis, S.J., Civin, C., and Small, D. (2012). FLT3-ITD knockin impairs hematopoietic stem cell quiescence/homeostasis, leading to myeloproliferative neoplasm. Cell Stem Cell *11*, 346–358.

Croce, C.M. (2008). Oncogenes and cancer. N. Engl. J. Med. 358, 502–511. DeKoter, R.P., and Singh, H. (2000). Regulation of B lymphocyte and macrophage development by graded expression of PU.1. Science 288, 1439–1441.

Ding, L., Ley, T.J., Larson, D.E., Miller, C.A., Koboldt, D.C., Welch, J.S., Ritchey, J.K., Young, M.A., Lamprecht, T., McLellan, M.D., et al. (2012). Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. Nature *481*, 506–510.

Enver, T., and Jacobsen, S.E. (2009). Developmental biology: Instructions writ in blood. Nature *461*, 183–184.

Gale, R.E., Green, C., Allen, C., Mead, A.J., Burnett, A.K., Hills, R.K., and Linch, D.C.; Medical Research Council Adult Leukaemia Working Party. (2008). The impact of FLT3 internal tandem duplication mutant level, number, size, and interaction with NPM1 mutations in a large cohort of young adult patients with acute myeloid leukemia. Blood *111*, 2776–2784.

Goardon, N., Marchi, E., Atzberger, A., Quek, L., Schuh, A., Soneji, S., Woll, P., Mead, A., Alford, K.A., Rout, R., et al. (2011). Coexistence of LMPP-like and GMP-like leukemia stem cells in acute myeloid leukemia. Cancer Cell *19*, 138–152.

Grossmann, V., Kern, W., Harbich, S., Alpermann, T., Jeromin, S., Schnittger, S., Haferlach, C., Haferlach, T., and Kohlmann, A. (2011). Prognostic relevance of RUNX1 mutations in T-cell acute lymphoblastic leukemia. Haematologica *96*, 1874–1877.

Growney, J.D., Shigematsu, H., Li, Z., Lee, B.H., Adelsperger, J., Rowan, R., Curley, D.P., Kutok, J.L., Akashi, K., Williams, I.R., et al. (2005). Loss of Runx1 perturbs adult hematopoiesis and is associated with a myeloproliferative phenotype. Blood *106*, 494–504.

Hegde, S., Ni, S., He, S., Yoon, D., Feng, G.S., Watowich, S.S., Paulson, R.F., and Hankey, P.A. (2009). Stat3 promotes the development of erythroleukemia by inducing Pu.1 expression and inhibiting erythroid differentiation. Oncogene *28*, 3349–3359.

Jacob, B., Osato, M., Yamashita, N., Wang, C.Q., Taniuchi, I., Littman, D.R., Asou, N., and Ito, Y. (2010). Stem cell exhaustion due to Runx1 deficiency is prevented by Evi5 activation in leukemogenesis. Blood *115*, 1610–1620.

Jan, M., Snyder, T.M., Corces-Zimmerman, M.R., Vyas, P., Weissman, I.L., Quake, S.R., and Majeti, R. (2012). Clonal evolution of preleukemic hematopoietic stem cells precedes human acute myeloid leukemia. Sci. Transl. Med. *4*, 149ra118.

Kharazi, S., Mead, A.J., Mansour, A., Hultquist, A., Böiers, C., Luc, S., Buza-Vidas, N., Ma, Z., Ferry, H., Atkinson, D., et al. (2011). Impact of gene dosage, loss of wild-type allele, and FLT3 ligand on Flt3-ITD-induced myeloproliferation. Blood *118*, 3613–3621.

Kiel, M.J., Yilmaz, O.H., Iwashita, T., Yilmaz, O.H., Terhorst, C., and Morrison, S.J. (2005). SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells. Cell *121*, 1109–1121.

Kundu, M., Compton, S., Garrett-Beal, L., Stacy, T., Starost, M.F., Eckhaus, M., Speck, N.A., and Liu, P.P. (2005). Runx1 deficiency predisposes mice to T-lymphoblastic lymphoma. Blood *106*, 3621–3624.

Lee, B.H., Tothova, Z., Levine, R.L., Anderson, K., Buza-Vidas, N., Cullen, D.E., McDowell, E.P., Adelsperger, J., Fröhling, S., Huntly, B.J., et al. (2007). FLT3 mutations confer enhanced proliferation and survival properties to multipotent progenitors in a murine model of chronic myelomonocytic leukemia. Cancer Cell *12*, 367–380.

Leow, S., Kham, S.K., Ariffin, H., Quah, T.C., and Yeoh, A.E. (2011). FLT3 mutation and expression did not adversely affect clinical outcome of childhood acute leukaemia—a study of 531 Southeast Asian children by the Ma-Spore study group. Hematol. Oncol. *29*, 211–219.

Levis, M., Murphy, K.M., Pham, R., Kim, K.T., Stine, A., Li, L., McNiece, I., Smith, B.D., and Small, D. (2005). Internal tandem duplications of the FLT3 gene are present in leukemia stem cells. Blood *106*, 673–680.

Li, L., Piloto, O., Nguyen, H.B., Greenberg, K., Takamiya, K., Racke, F., Huso, D., and Small, D. (2008). Knock-in of an internal tandem duplication mutation into murine FLT3 confers myeloproliferative disease in a mouse model. Blood *111*, 3849–3858.

Luc, S., Luis, T.C., Boukarabila, H., Macaulay, I.C., Buza-Vidas, N., Bouriez-Jones, T., Lutteropp, M., Woll, P.S., Loughran, S.J., Mead, A.J., et al. (2012). The earliest thymic T cell progenitors sustain B cell and myeloid lineage potential. Nat. Immunol. *13*, 412–419.

Månsson, R., Hultquist, A., Luc, S., Yang, L., Anderson, K., Kharazi, S., Al-Hashmi, S., Liuba, K., Thorén, L., Adolfsson, J., et al. (2007). Molecular evidence for hierarchical transcriptional lineage priming in fetal and adult stem cells and multipotent progenitors. Immunity *26*, 407–419.

Pecquet, C., Staerk, J., Chaligné, R., Goss, V., Lee, K.A., Zhang, X., Rush, J., Van Hees, J., Poirel, H.A., Scheiff, J.M., et al. (2010). Induction of myeloproliferative disorder and myelofibrosis by thrombopoietin receptor W515 mutants is mediated by cytosolic tyrosine 112 of the receptor. Blood *115*, 1037–1048.

Putz, G., Rosner, A., Nuesslein, I., Schmitz, N., and Buchholz, F. (2006). AML1 deletion in adult mice causes splenomegaly and lymphomas. Oncogene *25*, 929–939.

Rieger, M.A., Hoppe, P.S., Smejkal, B.M., Eitelhuber, A.C., and Schroeder, T. (2009). Hematopoietic cytokines can instruct lineage choice. Science *325*, 217–218.

Rumfelt, L.L., Zhou, Y., Rowley, B.M., Shinton, S.A., and Hardy, R.R. (2006). Lineage specification and plasticity in CD19- early B cell precursors. J. Exp. Med. *203*, 675–687.

Schmidt-Arras, D., Böhmer, S.A., Koch, S., Müller, J.P., Blei, L., Cornils, H., Bauer, R., Korasikha, S., Thiede, C., and Böhmer, F.D. (2009). Anchoring of FLT3 in the endoplasmic reticulum alters signaling quality. Blood *113*, 3568–3576.

Schnittger, S., Dicker, F., Kern, W., Wendland, N., Sundermann, J., Alpermann, T., Haferlach, C., and Haferlach, T. (2011). RUNX1 mutations are



frequent in de novo AML with noncomplex karyotype and confer an unfavorable prognosis. Blood *117*, 2348–2357.

Schwarz, B.A., Sambandam, A., Maillard, I., Harman, B.C., Love, P.E., and Bhandoola, A. (2007). Selective thymus settling regulated by cytokine and chemokine receptors. J. Immunol. *178*, 2008–2017.

Sitnicka, E., Bryder, D., Theilgaard-Mönch, K., Buza-Vidas, N., Adolfsson, J., and Jacobsen, S.E. (2002). Key role of flt3 ligand in regulation of the common lymphoid progenitor but not in maintenance of the hematopoietic stem cell pool. Immunity *17*, 463–472. Smith, C.C., Wang, Q., Chin, C.S., Salerno, S., Damon, L.E., Levis, M.J., Perl, A.E., Travers, K.J., Wang, S., Hunt, J.P., et al. (2012). Validation of ITD mutations in FLT3 as a therapeutic target in human acute myeloid leukaemia. Nature *485*, 260–263.

Steidl, U., Rosenbauer, F., Verhaak, R.G., Gu, X., Ebralidze, A., Otu, H.H., Klippel, S., Steidl, C., Bruns, I., Costa, D.B., et al. (2006). Essential role of Jun family transcription factors in PU.1 knockdown-induced leukemic stem cells. Nat. Genet. 38, 1269–1277.

Stirewalt, D.L., and Radich, J.P. (2003). The role of FLT3 in haematopoietic malignancies. Nat. Rev. Cancer 3, 650–665.