



# LUND UNIVERSITY

## **Blood-brain barrier permeability in rats exposed to electromagnetic fields used in wireless communication**

Persson, Bertil R; Salford, Leif G.; Brun, Arne

*Published in:*  
Wireless Networks

1997

[Link to publication](#)

*Citation for published version (APA):*

Persson, B. R., Salford, L. G., & Brun, A. (1997). Blood-brain barrier permeability in rats exposed to electromagnetic fields used in wireless communication. *Wireless Networks*, 3, 455-461.

*Total number of authors:*  
3

### **General rights**

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

# Blood-brain barrier permeability in rats exposed to electromagnetic fields used in wireless communication

Bertil R.R. Persson\*, Leif G. Salford and Arne Brun

Lund University, S-221 85 Lund, Sweden

Biological effects of *radio frequency* electromagnetic fields (EMF) on the blood-brain barrier (BBB) have been studied in Fischer 344 rats of both sexes. The rats were not anaesthetised during the exposure. All animals were sacrificed by perfusion–fixation of the brains under chloralhydrate anaesthesia after the exposure. The brains were perfused with saline for 3–4 minutes, and thereafter perfusion fixed with 4% formaldehyde for 5–6 minutes. Whole coronal sections of the brains were dehydrated and embedded in paraffin and sectioned at 5  $\mu\text{m}$ . Albumin and fibrinogen were demonstrated immunohistochemically and classified as normal versus pathological leakage. In the present investigation we exposed male and female Fischer 344 rats in a Transverse Electromagnetic Transmission line chamber to microwaves of 915 MHz as continuous wave (CW) and pulse-modulated with different pulse power and at various time intervals. The CW-pulse power varied from 0.001 W to 10 W and the exposure time from 2 min to 960 min. In each experiment we exposed 4–6 rats with 2–4 controls randomly placed in excited and non-excited TEM-cells respectively. We have in total investigated 630 exposed rats at various modulation frequencies and 372 controls. The frequency of pathological rats is significantly increased ( $p < 0.0001$ ) from 62/372 (ratio:  $0.17 \pm 0.02$ ) for control rats to 244/630 (ratio:  $0.39 \pm 0.03$ ) in all exposed rats. Grouping the exposed animals according to the level of specific absorbed energy (J/kg) give significant difference in all levels above 1.5 J/kg. The exposure was 915 MHz microwaves either pulse modulated (PW) at 217 Hz with 0.57 ms pulse width, at 50 Hz with 6.6 ms pulse width or continuous wave (CW). The frequency of pathological rats (0.17) among controls in the various groups is not significantly different. The frequency of pathological rats was 170/481 ( $0.35 \pm 0.03$ ) among rats exposed to pulse modulated (PW) and 74/149 ( $0.50 \pm 0.07$ ) among rats exposed to continuous wave exposure (CW). These results are both highly significantly different to their corresponding controls ( $p < 0.0001$ ) and the frequency of pathological rats after exposure to pulsed radiation (PW) is significantly less ( $p < 0.002$ ) than after exposure to continuous radiation (CW).

## 1. Introduction

The mammalian brain is protected from potentially harmful compounds in the blood by the so called blood-brain barrier (BBB). It is a selectively permeable, hydrophobic barrier that is readily crossed by small, lipid-soluble molecules [19,24]. Certain lipid-insoluble molecules such as glucose also readily crosses the cell layers constituting the barrier through carrier proteins that have a high affinity with specific molecules.

Although knowledge of the barrier's adaptive role is far from complete, there is a growing consensus that it serves not only to restrict entry of toxic polar molecules into the brain. It also serves as a regulatory system that stabilises and optimises the fluid environment of the brain's intracellular compartment [19,24].

The intact BBB protects the brain from damage, whereas a dysfunctioning BBB, as induced by epileptic seizures or extreme hypertension, allows influx of normally excluded hydrophilic molecules into the brain tissue. This might lead to cerebral oedema, increased intracranial pressure and in the worst case, irreversible brain damage.

The normal selective permeability of the blood-brain barrier (BBB) can be altered in several neuropathological

conditions and experimental situations. The BBB opens during an acute hypertensive episode [30], in epileptic seizures [15,31], and during therapeutic X-ray irradiation [16].

Several authors have reported that electromagnetic exposure alters BBB permeability [1–4,7,8,10–12,17,20,21,32,33]. Other authors have reported difficulties in confirming these findings [11,14,22,23,35–38]. The lack of agreement among investigators might be due to lack of an assay of BBB permeability that combines sensitivity and specificity.

Shivers et al. [29] observed during exposure to clinical magnetic resonance imaging (MRI) procedures an amplified vesicle-mediated transport of horseradish peroxidase across the endothelium to the extracellular compartment of the brain parenchyma of the rat. They did not, however, report on effects of the exclusive field components. The mechanism involved, however, points strongly to facilitation of *pinocytosis*-like transport of albumin through BBB [18].

In the present investigation we have studied the permeability of BBB to endogenous albumin and fibrinogen during exposure to microwaves. We exposed rats to 915 MHz microwaves as continuous wave (CW) and pulse-modulated at the various repetition rates ( $4\text{--}217\text{ s}^{-1}$ ). Preliminary results are previously reported by Salford et al. [25–28].

Our model does not make use of radioactive tracers and does not supply foreign substances or anaesthetics to the

\* Corresponding author: Radiation Physics Department, Lund University Hospital, S-221 85 Lund, Sweden.  
E-mail: Bertil.Persson@radfys.lu.se

rat during exposure. Instead it uses the animals own albumin, that does not penetrate the intact BBB. The passage of even minute amounts of albumin from the capillaries through an opened BBB into the surrounding brain tissue is revealed by our histopathological technique. This also allows for the identification of exactly in which constituents (neurones, glial cells, extracellular space) of the surrounding brain, where the albumin is situated. The perfusion–fixation method washes out all blood and its albumin content from the brain vessels including the capillaries. This facilitates the identification of albumin that has leaked out through the opened BBB into the brain.

## 2. Material and methods

### 2.1. Exposure in a TEM-cell

A Transverse Electromagnetic transmission line cell (TEM-cell) used for the RF exposure of rats was designed by dimensional scaling from previously constructed cells at the National Bureau of Standards (Crawford, 1974). TEM-cells are known to generate uniform TEM-fields for standard measurements.

The cell is enclosed in a wooden box that supports the outer conductor and central plate. The outer conductor is made of brass-net and is attached to the inner walls of the box. The centre plate, or septum, is constructed of aluminium and is held up by teflon braces which are screwed at the inner side walls.

To allow access to the inside of the cell both ends can be removed. The inside of the cell is ventilated through 18 holes (diam. 18 mm) in the side walls and top of the box and the brass-net of 50 mesh allows air to circulate. These holes are also used for examination of the interior during exposure. Probes for monitoring temperature inside the cell or of test object are inserted through these holes.

The rats are placed in plastic trays to avoid contact with the central plate and outer conductor. The bottom of the tray is covered with absorbing paper to collect urine and faeces.

The rats were exposed to 915 MHz electromagnetic radiation continuous wave and pulse-modulated with different repetition rates. The modulated RF-radiation consists of square wave shaped pulses with durations of 0.57, 4 or 6 ms and intensities (in Watts) during the presence of the pulse. Transmitted and absorbed power was measured at continuous wave exposure with and without rats in the TEM-cell. From these measurements the average SAR in the whole rat was calculated to be  $1.2 \pm 0.4$  W/kg per watt of input power. This value was in good agreement with the theoretical estimate of 1.6 W/Kg per watt of input power that was used in the evaluation of the experiment [5,9,13,34].

### 2.2. Albumin and fibrinogen immunohistochemistry

Fischer 344 rats of both sexes, weighing 119–555 g (median: 202 g; 25% quartiles: 175 g; 75% quartiles: 273 g)

were used in these experiments (own breeding). The rats were not anaesthetised, during the exposure.

Both controls and exposed animals were sacrificed by perfusion–fixation of the brains under chloralhydrate anaesthesia between 20 minutes and 2 hours after the exposure. The brains were perfused with saline for 3–4 minutes, thereafter fixed in 4% formaldehyde for 5–6 minutes and immersion fixed in 4% formaldehyde for more than 24 hours. Whole coronal sections of the brains (3, 7 and 11 mm from the tip of the frontal pole) were dehydrated and embedded in paraffin and sectioned at 5  $\mu$ m. The chloralhydrate anaesthesia is necessary to avoid stress and blood pressure rise during perfusion–fixation procedure. Also for ethical reason no animals were sacrificed without chloralhydrate anaesthesia.

Albumin was demonstrated with the IgG fraction of rabbit anti-rat albumin (Cappel Research Products, Organon Teknika, Västra Frölunda, Sweden) diluted 1:16,000. Fibrinogen was demonstrated with rabbit anti-human fibrinogen (Dacopatts AB, Hägersten, Sweden), diluted 1:500. Incubation time for both was over night at 4°C.

Biotinylated swine anti-rabbit IgG was used as a secondary antibody. Then avidin, peroxidase conjugated, was coupled to the biotin and visualised with DAB (diaminobenzidine), counterstained with Meyer-HTX [6]. Standard control procedures were used for both albumin and fibrinogen.

The numbers of immunopositive extravasates were recorded under a microscope. None or occasional minor leakage was rated as normal, whereas one larger or several leakages were regarded as pathological. Immunopositive sites were, however, disregarded when localised in the hypothalamus, basally from the median eminence and laterally including the nucleus lateralis hypothalami, in the immediate vicinity of the third ventricles. These structures are well known for their insufficient blood-brain barrier and within any part of the choroid plexus of the ventricles consistently shows immunopositivities, mostly of a diffuse type, in the strain used in the present experiments.

### 2.3. Statistics

The frequency of occurrence of albumin extravasation in exposed and control animals were compared with chi-square or Fisher's exact probability test.

## 3. Results and discussions

Example of pathological leakage around small vessels is demonstrated in figure 1. The number of pathological rat brains among all control rats is 62 out of 372 (ratio:  $0.17 \pm 0.02$ ). These findings are occasional and rare and are probably due to normal minor disturbances. The frequency of pathological rats among controls in the various groups is not significantly different ( $p < 0.4$ ).

In the present investigation we exposed male and female Fischer 344 rats in a TEM-chamber to microwaves of 915 MHz as continuous wave (CW) and pulse-modulated




Figure 1. Example of pathological leakage around small vessels demonstrated through immunostaining against albumin. Hematoxylin eosin staining. (a) Section through frontal lobe with cortex and white matter, magnification  $\times 5$ . (b) Part of (a) with vessel, magnification  $\times 10$ .

with different pulse power and at various time intervals. The CW-pulse power varied from 0.001 W to 10 W and the exposure time from 2 min to 960 min. In each experiment we exposed 4 rats with 4 controls randomly placed in excited and non-excited TEM-cells, respectively.

We have in total investigated 630 exposed rats at various modulation frequencies and 372 controls. The frequency of pathological rats is significantly increased ( $p < 0.0001$ ) from 62/372 (ratio:  $0.17 \pm 0.02$ ) for control rats to 244/630 (ratio:  $0.39 \pm 0.03$ ) in all exposed rats. Grouping the animals according to the level of specific absorbed energy (J/kg) that is normally referred to as "Specific Absorption" (SA), give significant difference ( $p < 0.0001$ ) in all levels of absorbed energy above 1.5 J/kg compared to controls. (see table 1).

The exposure was 915 MHz microwaves either pulse modulated (PW) at 217 Hz with 0.57 ms pulse width, at 50 Hz with 6.6 ms pulse width or continuous wave (CW). The frequency of pathological rats ( $< 0.2$ ) among controls in the various groups is not significantly different ( $p < 0.4$ ). The frequency of pathological rats was 170/481 ( $0.35 \pm 0.03$ ) among rats exposed to pulse modulated (PW) and 74/149 ( $0.50 \pm 0.07$ ) among rats exposed to continuous wave exposure (CW). These results are both

highly significantly different to their corresponding controls ( $p < 0.0001$ ) and the frequency of pathological rats after exposure to pulsed radiation (PW) is significantly less ( $p < 0.002$ ) than after exposure to continuous radiation (CW) (table 2). The degree of pathological leakage in exposed animals is more severe and more frequent per animal compared to the controls.

The results of BBB-permeability of albumin in rats exposed to 915 MHz microwaves with different modulation frequencies for groups exposed to similar SAR values are displayed in figures 2–5. In figure 2 are displayed the results at SAR values  $4 \times 10^{-4}$ – $8 \times 10^{-3}$  W/kg. Although the SAR values are very low, around 1 mW/kg, this group indicates the highest fraction of pathological findings recorded in the entire investigation. There seems to be a maximum effect around 8–50 Hz modulation frequency. In figure 3 is given the results at SAR values  $(2-8) \times 10^{-2}$  W/kg. In this group the effect of CW and 50 Hz modulation is the same but the effect of 16 Hz and 8.3 & 217 Hz GSM modulation is very low and not significant. The overall effect of pulse modulated exposure is, however, significant. Figure 4 shows the results at SAR values 0.11–0.95 W/kg. The effect of CW exposure is about the same as in previous group but the effect of 217 Hz pulse modulation shows

Table 1  
BBB-permeability of rat brain after 915 MHz RF-exposure at various intervals of specific absorption SA (J/kg).

Group of specific absorption (J/kg)	Normal score < 1	Pathological score ≥ 1	Ratio ± SD Pathological/all	Chi-square # Contr. p
Controls SA = 0	310	62	0.17 ± 0.02	
1.5 ≤ SA < 10	32	34	0.52 ± 0.11	< 0.00005
10 ≤ SA < 100	30	20	0.40 ± 0.11	< 0.0002
100 ≤ SA < 1000	116	61	0.34 ± 0.05	< 0.00005
1000 ≤ SA < 10 000	122	78	0.39 ± 0.05	< 0.00005
10 000 ≤ SA < 48 000	90	52	0.37 ± 0.01	< 0.00005
All exposed (630)	386	244	0.39 ± 0.03	< 0.00005

Table 2  
BBB-permeability of rat brain after exposure to 915 MHz microwaves either Pulsed (PW) or Continuous Wave (CW).

Group	Number of rats with normal score < 1	Number of rats with pathological score ≥ 1	Ratio ± SD No. score ≥ 1 to all	p Chi-square exposed versus controls	p Chi-square PW versus CW
Controls pulsed (259)	219	40	0.15 ± 0.03	0.68	0.34
Controls CW (113)	91	22	0.19 ± 0.05	0.49	
All controls (372)	310	62	0.17 ± 0.02		
All exposed pulsed (481)	311	170	0.35 ± 0.03	< 0.00005	< 0.002
All exposed CW (149)	75	74	0.50 ± 0.07	< 0.00005	
All exposed (630)	386	244	0.39 ± 0.03	< 0.00005	




Figure 2. Blood-brain barrier permeability of albumin in rats exposed to 915 MHz microwaves with different modulation frequencies at SAR values  $4 \times 10^{-4}$ – $8 \times 10^{-3}$  W/kg. The average fraction of pathological leakage in 372 controls was  $0.17 \pm 0.02$ .

a significant effect compared to controls. The effect of 8.3 and 16 Hz modulation is however not significantly different from the controls. Figure 5 displays the results at SAR values 1.7–8.3 W/kg. In this group the SAR levels approach the threshold for thermal effects and the effect of CW exposure is very pronounced with a fraction of pathological rats of 55%. Surprisingly the effect of pulse modulated exposure is not significant at this high SAR level.

The frequency of pathological rats was 170/481 ( $0.35 \pm 0.03$ ) among rats exposed to pulse modulated (PW) and 74/149 ( $0.50 \pm 0.07$ ) among rats exposed to continu-




Figure 3. Blood-brain barrier permeability of albumin in rats exposed to 915 MHz microwaves with different modulation frequencies at SAR values  $(2-8) \times 10^{-2}$  W/kg. The average fraction of pathological leakage in 372 controls was  $0.17 \pm 0.02$ .

ous wave exposure (CW). These results are both highly significantly different from their corresponding controls ( $p < 0.0001$ ) and the frequency of pathological rats after exposure to pulsed radiation (PW) is significantly less ( $p < 0.002$ ) than after exposure to continuous radiation (CW) (table 2). This is a highly interesting observation as the current opinion is that pulse modulated electromagnetic fields are more potent in causing biological effects.

We have demonstrated that microwave exposure produces an unequivocal effect on the BBB in our rats. The clinical importance of this finding, however, is disputable. Our method for detection of albumin is extremely sensi-




Figure 4. Blood-brain barrier permeability of albumin in rats exposed to 915 MHz microwaves with different modulation frequencies at SAR values 0.11–0.95 W/kg. The average fraction of pathological leakage in 372 controls was  $0.17 \pm 0.02$ .




Figure 5. Blood-brain barrier permeability of albumin in rats exposed to 915 MHz microwaves with different modulation frequencies at SAR values 1.7–8.3 W/kg. The average fraction of pathological leakage in 372 controls was  $0.17 \pm 0.02$ .

tive and reveals even minute amounts of albumin leaking through the BBB, so small that they may be harmless to the brain. However, the potential health hazards of the opening the BBB during exposure to wireless communication demands further investigation.

## Acknowledgements

The authors thank Kerstin Stureson and Ylva Knutsson who helped with the histopathological preparations, Susanne Strömblad and Catharina Blennow who handled the animals with great skill, and Lars Malmgren for technical support.

## References

[1] E.N. Albert, Light and electron microscopic observation on the blood-brain barrier after microwave irradiation, in: *Symp. on Biol. Eff. and Measur. of Radiofr/Microwaves, FDA 77-8026*, ed. D.G. Hazzard (HEW Publications, Washington, DC, 1977) pp. 294–304.

[2] E.N. Albert, D.L. Brainard, J.D. Randall and F.S. Janatta, Neuro-pathological observations of microwave irradiated hamsters, in: *URSI Int. Symp. Biol. Eff. Electromagn. Radiat.* (URSI, Helsinki, 1978) p. 58.

[3] E.N. Albert and S. Mahmoud, Morphological changes in cerebellum of neonatal rats exposed to 2.45 GHz microwaves, in: *Electromagnetic Fields and Neurobehavioral Function*, eds. M.E. O'Connors and R.H. Lovely (Alan R. Liss, Inc., New York, 1984) pp. 135–152.

[4] Y. Ashani, F.H. Henry and G.N. Catravas, Combined effects of anti-cholinesterase drugs and low-level microwave radiation, *Radiat. Res.* 84 (1980) 496–503.

[5] M.L. Crawford, Generation of standard EM field using TEM transmission cells, *IEEE Trans. Electromagn. Compat.* 16 (1974) 189–195.

[6] Dacopatt, *Handbook of Immunochemical Staining Methods* (1994).

[7] H.J. Garber, W.H. Oldendorf, L.K. Brown and R.B. Lufkin, MRI gradient fields increase brain mannitol space, *Magn. Reson. Imaging* 7 (1989) 605–610.

[8] H. Goldman, J.C. Lin, S. Murphy and M.F. Lin, Cerebral permeability of  $^{86}\text{Rb}$  in the rat after exposure to pulsed microwaves, *Bioelectromagnetics* 5 (1984) 323–330.

[9] J.W. Hand, Biophysics and technology of electromagnetic hyperthermia, in: *Methods of External Hyperthermia Heating*, ed. M. Gauthier (Springer, Berlin, 1990) pp. 1–59.

[10] J.C. Lin, Engineering and biophysical aspects of microwave and radiofrequency radiation, in: *Hyperthermia*, eds. D.J. Watmough and W.M. Ross (Blackie and Son, Glasgow, 1986) pp. 42–75.

[11] J.C. Lin and M.F. Lin, Studies on microwave and blood-brain barrier interaction, *Bioelectromagnetics* 1 (1980) 313–323.

[12] J.C. Lin and M.F. Lin, Microwave hyperthermia-induced blood-brain barrier alterations, *Radiat. Res.* 89 (1982) 77–87.

[13] L. Martens, J. Van Hese, D. De Zutter, C. De Wagter, L. Malmgren, B. Persson and L.G. Salford, Electromagnetic field calculations used for exposure experiments on small animals in TEM-cells, *Bioelectrochemistry and Bioenergetics* 30 (1992) 313–318.

[14] J.H. Merritt, A.P. Chamness and S.J. Allen, Studies on blood-brain barrier permeability after microwave radiation, *Radiat. Environ. Biophys.* 15 (1978) 367–377.

[15] A. Mihaly and B. Bozoky, Immunohistochemical localization of serum proteins in the hippocampus of human subjects with partial and generalized epilepsy and epileptiform convulsions, *Acta Neuropathol.* 127 (1984) 251–267.

[16] V. Nair and L.J. Roth, Effect of X-irradiation and certain other treatments on blood-brain barrier permeability, *Radiat. Res.* 23 (1964) 249–269.

[17] J.P. Neilly and J.C. Lin, Interaction of microwaves on the blood-brain barrier of rats, *Bioelectromagnetics* 7 (1986) 405–414.

[18] C. Neubauer, A.M. Phelan, H. Kues and D.G. Lange, Microwave irradiation of rats at 2.45 GHz activates pinocytotic-like uptake of tracer by capillary endothelial cells of cerebral cortex, *Bioelectromagnetics* 11 (1990) 261–268.

[19] W.H. Oldendorf, Permeability of the blood-brain barrier, in: *The Nervous System*, ed. D. Tower (Raven Press, New York, 1975) pp. 229–289.

[20] K.J. Oscar and T.D. Hawkins, Microwave alteration of the blood-brain barrier system of rats, *Brain Res.* 126 (1977) 281–293.

[21] F.S. Prato, J.R.H. Frappier, R.R. Shivers, M. Kavaliers, P. Zabel, D.J. Drost and T.Y. Lee, Magnetic resonance imaging increases the blood-brain barrier permeability to 153-gadolinium diethylenetriaminepentaacetic acid in rats, *Brain Res.* 523 (1990) 301–304.

[22] E. Preston and G. Prefontaine, Cerebrovascular permeability to sucrose in the rat exposed to 2450 MHz microwaves, *Appl. Physiol. Respir. Environ. Exercise Physiol.* 49 (1980) 218–223.

[23] E. Preston, E.J. Vavasour and H.M. Assenheim, Permeability of the blood-brain barrier to mannitol in the rat following 2,450 MHz microwave irradiation, *Brain Res.* 174 (1979) 109–117.

[24] S.I. Rapoport, *Blood-Brain Barrier in Physiology and Medicine* (Raven Press, New York, 1976).

[25] L.G. Salford, A. Brun, J. Eberhardt, L. Malmgren and B. Persson,


- Electromagnetic field-induced permeability of the blood-brain barrier shown by immunohistochemical methods, in: *Interaction Mechanism of Low-Level Electromagnetic Fields in Living Systems*, eds. B. Nordén and C. Ramel (Oxford University Press, Oxford, 1992) pp. 251–258.
- [26] L.G. Salford, A. Brun, J.L. Eberhardt and B.R.R. Persson, Permeability of the blood-brain barrier induced by 915 MHz electromagnetic radiation, continuous wave and modulated at 8, 16, 50, and 200 Hz, *Biochemistry and Bioenergetics* 30 (1993) 293–301.
- [27] L.G. Salford, A. Brun, J.L. Eberhardt and B.R.R. Persson, Permeability of the blood/brain barrier induced by 915 MHz electromagnetic radiation, CW and modulated at various SARs, in: *Electricity and Magnetism in Biology and Medicine*, ed. M. Blank (San Francisco Press Inc., San Francisco, CA, 1993).
- [28] L.G. Salford, A. Brun, K. Sturesson, J.L. Eberhardt and B.R.R. Persson, Permeability of the blood-brain barrier induced by 915 MHz electromagnetic radiation, continuous wave and modulated at 8, 16, 50, and 200 Hz, *Microscopy Research and Technique* 27 (1994) 535–542.
- [29] R.R. Shivers, M. Kavaliers, G.C. Teskey, F.S. Prato and R.M. Pelletier, Magnetic resonance imaging temporarily alters blood-brain barrier in the rat, *Neuroscience Letters* 76 (1987) 25–31.
- [30] T.-E.O. Sokrab, B.B. Johansson, H. Kalimo and Y. Olsson, A transient hypertensive opening of the blood-brain barrier can lead to brain damage, *Acta Neuropathology* 75 (1988) 557–565.
- [31] T.-E.O. Sokrab, H. Kalimo and B.B. Johansson, Parenchymal changes related to plasma protein extravasation in experimental seizures, *Epilepsia* 31 (1990) 1–8.
- [32] C.H. Sutton and F.B. Carrol, Effects of microwave-induced hyperthermia on the blood-brain barrier of the rat, *Radiat. Sci.* 14 (1979) 329–334.
- [33] C.H. Sutton, R.L. Nunnally and F.B. Carroll, Protection of the microwave-irradiated brain with body core hypothermia, *Cryobiology* 10 (1973) 513–514.
- [34] J. Van Hese, L. Martens, D. De Zutter, C. De Wagter, L. Malmgren, B.R.R. Persson and L.G. Salford, Simulation of the effect of inhomogenities in TEM transmission cells using the FDTD-method, *IEEE Trans. Electromagn. Compat.* 34 (1991) 292–298.
- [35] W.M. Williams, M. del Cerro and S.M. Michaelson, Effect of 2450 MHz microwave energy on the blood-brain barrier to hydrophilic molecules. B. Effect on permeability to HPR, *Brain Res. Rev.* 7 (1984) 171–182.
- [36] W.M. Williams, W. Hoss, M. Formaniak and S.M. Michaelson, Effect of 2450 MHz microwave energy on the blood-brain barrier to hydrophobic molecules. A. Effect on permeability to sodium fluorescein, *Brain Res. Rev.* 7 (1984) 165–170.
- [37] W.M. Williams, S.-T. Lu, M. del Cerro and S.M. Michaelson, Effect of 2450 MHz microwave energy on the blood-brain barrier to hydrophilic molecules. D. Brain temperature and blood-brain barrier permeability to hydrophilic molecules, *Brain Res. Rev.* 7 (1984) 191–212.
- [38] W.M. Williams, J. Platner and S.M. Michaelson, Effect of 2450 MHz microwave energy on blood-brain barrier to hydrophilic molecules. C. Effect on permeability to [<sup>14</sup>C]sucrose, *Brain Res. Rev.* 7 (1984) 183–190.



**Bertil R.R. Persson**, Ph.D., professor of medical radiation physics, was born 1938 in Malmö, Sweden, studied at University of Lund, Sweden, and in 1970 became doctor of philosophy and associate professor in radiation physics. From July 1980 till present he is full professor in medical radiation physics at Lund University hospital where he is head of the department of radiation physics. During his scientific activity he has published more than 320 scientific publications and written 16 extensive reports and books. His scientific career began in 1963 with studies

of the fall-out from atmospheric nuclear weapons tests in the food-chain lichen-reindeer-man and continued in medical use of short-lived radioisotopes. The scientific activity in environmental radiology has developed into polar research. In 1980 he participated in the Swedish Arctic Expedition Ymer-80, in January–April 1989 he participated in the Swedish Antarctic research program with a marine ecology program investigating the longitudinal profile of <sup>134</sup>Cs/<sup>137</sup>Cs ratio from Gothenburg in Sweden all the way to Antarctica. In June–August 1994 he participated in the Swedish–Russian Tundra expedition with marine and terrestrial radioecology research along the coast of Siberia and in 1995 he studied the environmental radioactivity in and around the Russian uranium mines in Krasnokamensk. He just returned from the Swedish expedition Arctic Ocean-96 that visited the North Pole 1996-09-10 the latest visit ever. During the expedition he studied the oceanic transport of radioactive elements, UV-radiation and cosmic rays (myons).


In 1977 he began to study microwave induced hyperthermia for tumour-therapy and since then this project has expanded very rapidly. He developed an equipment for local hyperthermia in patients with breast carcinoma, and recently equipment for heat treatment of benign prostate hyperplasia and menorrhagia. He started already in 1982 with biomedical applications of nuclear magnetic resonance (NMR) and has developed and built a special equipment for performing *in vivo* NMR studies in both animal and man. In close co-operation with several other scientists he is also conducting *in vitro* studies of the NMR relaxation times in tissues of thyroid and brain, also studying the influence of paramagnetic ions and super-paramagnetic particles. He has studied extensively clinical measurements of flow, microcirculation, diffusion and perfusion with NMR. During the past years he has been deeply involved in studying the effects of electromagnetic fields on healthy brain and on brain-tumours. He was co-sponsor for the New York Academy of Science conference on the biological and health effects of clinical NMR examination held in 1991. At present he is deeply involved in studies of the effect of low power pulsed microwave fields on the blood brain barrier and other biological systems. He is also engaged in using electromagnetic fields for tumour therapy and has made a lot of progress in the use of high voltage electric fields for permeabilization of tumour cell membranes *in vivo*. The objective of that investigation is to study the destructive effect of electropermeabilization on tumours and its ability to introduce toxic compounds into tumour cells. He is studying the therapeutic effect of electropermeabilization on tumour cells in presence of chemotherapeutic agents (for example, Bleomycin) and radioimmunotherapeutic agents.



**Leif G. Salford** was born in Malmö, Sweden, in 1941. He received the MD degree from the University of Lund in 1969, and the Ph.D. degree from the same university in 1974 with the thesis "Influence of profound hypoxia on regional metabolism, blood flow and cell morphology in rat brain". During 1972–1973 he was Wrightsman scholar at the Dept. of Neurology, Cornell Medical Center, New York Hospital, NY. In 1977 he became associate professor of neurosurgery, Lund University, and in 1979 Consultant Neurosurgeon. He was Professor and Chairman, Department of Neurosurgery, Kuwait University, 1981–1983, and Professor of Neurosurgery and Director of the Departments of Clinical Neurosciences, Sahlgrenska University Hospital and Göteborg University, Sweden, 1993–1996. His actual positions are: Professor and Chairman, Dept. of Neurosurgery, Lund University Hospital, and Director of the Institute for Clinical Neuroscience, Lund University, Lund, Sweden. Dr. Salford is Chairman of the Neuro-oncology Committee of the World Federation of Neurosurgical Societies, and President of the European Association for Neuro-oncology. He is President of the Swedish Neurosurgical Society, Honorary President of the Scandinavian Neurooncology Society and Expert in the EU Information Society Forum.

Dr. Salford's research is concentrated on the malignant primary brain tumours and on the search for efficient treatment against this hitherto incurable type of cancer. This has also resulted in his studies of opening the

blood-brain-barrier by the use of electromagnetic fields in order to reach the tumour cells with cytotoxins, and in his interest in the biophysical effects of EMF upon brain and tumour biology. Dr. Salford is Director of the Laboratory for Experimental Neurooncology, Lund University, where animal models and tissue cultures play an important role in his research in close cooperation with technological institutions such as Medical Radiation Physics. He strives to utilize the latest progress in technology in his search for solutions to medical problems. Another result of this collaboration is the introduction of electroporation *in vivo* as a tool for treatment of intraparenchymatous tumours which has proven efficient in connection with cytotoxins in the rodent brain tumour model.



**Arne Brun**, MD, Ph.D., is a professor of neuropathology, University Hospital, Lund, Sweden. His publications are mainly on central nervous system diseases such as dementias, especially the Alzheimers' disease and frontal dementias, also cerebrovascular incl. BBB-studies.