



LUND UNIVERSITY

Palaeokarst formation in the early Palaeozoic of Baltoscandia : evidence for significant sea-level changes in a shallow epicontinental sea

Lehnert, Oliver; Calner, Mikael; Ahlberg, Per; Ebbestad, Jan Ove; Harper, David A.T.; Meinhold, Guido

Published in:
[Publication information missing]

2013

[Link to publication](#)

Citation for published version (APA):

Lehnert, O., Calner, M., Ahlberg, P., Ebbestad, J. O., Harper, D. A. T., & Meinhold, G. (2013). Palaeokarst formation in the early Palaeozoic of Baltoscandia : evidence for significant sea-level changes in a shallow epicontinental sea. *[Publication information missing]*, 169-171.

Total number of authors:
6

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Proceedings of the 3rd IGCP 591 Annual Meeting

Lund, Sweden, 9–19 June 2013



Edited by

Anders Lindskog and Kristina Mehlqvist

Suggested reference format

Lindskog, A. & Mehlqvist, K., 2013: *Proceedings of the 3rd IGCP 591 Annual Meeting – Lund, Sweden, 9–19 June 2013*. Lund University. 368 pp.

Hinnov, L.A., 2013: Prospects for a Paleozoic Astronomical Time Scale. In A. Lindskog & K. Mehlqvist (eds.): *Proceedings of the 3rd IGCP 591 Annual Meeting – Lund, Sweden, 9–19 June 2013*, 19–20. Lund University.

Department of Geology
Lund University
Sölvegatan 12
SE-223 62 Lund
Sweden

<http://www.geol.lu.se>

<http://www.igcp591.org>

ISBN: 978-91-86746-87-2



Foreword and acknowledgements

This abstract volume has been prepared for the 3rd annual meeting of the IUGS/UNESCO International Geoscience Programme Project 591 *The Early to Middle Palaeozoic Revolution*. The meeting was hosted by the Department of Geology, Lund University, in June 9–19 2013 and followed the successful annual meetings held in Madrid/Ludlow (2011) and Cincinnati (2012). The Lund conference was arranged jointly with the annual meetings of the Cambrian, Ordovician and Silurian subcommissions on stratigraphy, and included a post-conference excursion to key geological localities in Skåne, Västergötland and the Oslo Region. The conference was a focus for cutting-edge research in Lower and Middle Palaeozoic geology and palaeontology, and the presentations covered a wide range of topics from morphology and taxonomy of various fossil groups through advances in geochemistry and stratigraphy to biogeography, palaeoecology and palaeoclimatology. We would like to express our sincere gratitude to Anders Lindskog and Kristina Mehlqvist for their meticulous editing of the meeting proceedings. We are also grateful for valuable input from the organization and scientific committee associated with the meeting. We acknowledge financial support from the Swedish Research Council (grant D0013001 to MC), the Geological Survey of Sweden, the Geological Society of Sweden, the Department of Geology at Lund University, and the municipality of Lund.

Lund on 8 May 2013

Mikael Calner (meeting chair)

Oliver Lehnert (vice chair)

Per Ahlberg

Palaeokarst formation in the early Palaeozoic of Baltoscandia – evidence for significant sea-level changes in a shallow epicontinental sea

OLIVER LEHNERT^{1,2}, MIKAEL CALNER², PER AHLBERG², JAN OVE EBBESTAD³, DAVID A. T. HARPER⁴ AND GUIDO MEINHOLD⁵

In the Lower Palaeozoic sedimentary succession of Sweden palaeokarsts have been reported from different stratigraphic levels in the Silurian strata of Gotland by Calner (2008; see references therein). Until last year there were no records of Cambrian karsts and in the Ordovician only the basin-wide Katian palaeokarst horizon in the Upper Ordovician Slandrom Limestone has been described in detail (Calner et al. 2010a). The unconformities and disconformities on top of the slightly older Kullsberg mounds in quarries located in the Siljan impact structure (Dalarna) presumably represent an earlier regression and karstic development (Calner et al. 2010b). Beside these reports, there is only the statement by Nielsen (1995) that karst may have formed at the top of the Darriwilian Komstad Limestone. During the last two years, however, several new and significant palaeokarst surfaces have been detected in the Cambrian–Ordovician successions of Sweden (Lehnert et al. 2012).

At Kakeled Quarry (Västergötland), a palaeokarst cave with a breccia fill (large, angular Orsten clasts in a dark limestone matrix) is exposed beneath a ‘Middle Cambrian’ palaeokarst surface (Jiangshanian Stage) located close to the top of the Kakeled Limestone Bed of the Alum Shale Formation (Lehnert et al., 2012). In the karstic pockets, a mass occurrence of *Orusia lenticularis* occurs. These shallow-water brachiopods originally settled on hard substrates after a major regression exposing, regionally, the sea floors of the alum shale basin. Their reworking and concentration in the conglomeratic bed overlying the irregular palaeokarst surface reflects deposition during transgression in extremely shallow marine environments.

A younger karst surface is exposed in Tomten Quarry at Torbjörntorp (Västergötland). In two dimensions in the quarry wall it resembles the “Schrattenkalk”, but rock slabs cut vertically and parallel to bedding planes display a karren system that resembles “Napfkarren” or cockling features. Trilobites of the Furongian *Ctenopyge bisulcata* and *C. linnarssoni* zones occur in the 1–2 cm thick, glauconitic packstone bed that overlies the palaeokarst surface and which represents the upper Tremadocian Björkåsholmen Formation. The associated stratigraphic gap comprises the six uppermost trilobite zones of the Furongian plus most of the Tremadocian. Darriwilian conodonts with reworked

older material within a limestone bed slightly above the glauconitic packstone bed indicate yet another substantial gap in the succession.

In the new Tingskullen core from northeastern Öland, another palaeokarst surface with grikes and evidence of repeated exposure marks the top of the upper Tremadocian *Obolus* conglomerate (?) or a lower limestone part of the Djupvik Formation (“*Ceratopyge* Shale”). This palaeokarst surface is overlain by glauconitic limestone of the Köpingsklint Formation and inferably reflects the global *Ceratopyge* Regressive Event (CRE).

At the base of the Lanna Limestone in the Siljan area, palaeokarst is associated with the Dapingian Blommiga Bladet (‘flowery sheet’) hardground complex, which can be correlated across most of Baltoscandia.

The basin-wide palaeokarst in the Katian Slandrom Limestone (Calner et al. 2010a) no longer marks the youngest Ordovician karst record. Recently, Hirnantian karst caves and solution cavities filled with greenish marls of the Glisstjärn Formation have been recognized in sections of the Boda Limestone in the Siljan Ring structure (Dalarna). Solution and karst cave formation reflects an interval of the regression during the Hirnantian glaciation and the youngest period of subaerial exposure during the Ordovician.

Some earlier sedimentary models suggesting that Baltoscandia was flooded by a deep epicontinental sea are challenged by the discovery of multiple palaeokarst development together with other shallow-water features. Instead, palaeokarst formation implies subaerial exposure during a number of major regressions.

¹*GeoZentrum Nordbayern, Lithosphere Dynamics, University of Erlangen-Nürnberg, Schloßgarten 5, D-91054, Erlangen, Germany; lehnert@geol.uni-erlangen.de*

²*Department of Geology, Lund University, Sölvegatan 12, SE-223 62 Lund, Sweden*

³*Museum of Evolution, Uppsala University, Norbyvägen 16, SE-752 36 Uppsala, Sweden*

⁴*Department of Earth Sciences, Durham University, Durham DH1 3LE, UK*

⁵*Geowissenschaftliches Zentrum der Universität Göttingen, Abteilung Sedimentologie/Umweltgeologie, Goldschmidtstraße 3, D-37077 Göttingen, Germany*

References

- Calner, M., 2008: Silurian global events – at the tipping point of climate change. In A.M.T. Elewa (ed.): *Mass extinction*, 21–57. Springer Book.
- Calner, M., Lehnert, O. & Nölvak, J., 2010a: Palaeokarst evidence for widespread regression and subaerial exposure in the middle Katian (Upper Ordovician) of Baltoscandia: Significance for global climate. *Palaeogeography, Palaeoclimatology, Palaeoecology* 296, 235–247.
- Calner, M., Lehnert, O. & Joachimski, M. 2010b: Carbonate mud mounds, conglomerates, and sea-level history in the Katian (Upper Ordovician) of central Sweden. *Facies* 56, 157–172.

- Lehnert, O., Calner, M., Ahlberg, P. & Harper, D.A. 2012: Multiple palaeokarst horizons in the Lower Palaeozoic of Baltoscandia challenging the dogma of a deep epicontinental sea. *Geophysical Research Abstracts* 14, EGU2012-11362-1.
- Nielsen, A.T., 1995: Trilobite systematics, biostratigraphy and palaeoecology of the Lower Ordovician Komstad Limestone and Huk formations, southern Scandinavia. *Fossils and Strata* 38, 1–374.