



LUND UNIVERSITY

Photolysis-free two photon laser induced fluorescence of H atom

Ruchkina, Maria; Ding, Pengji; Aldén, Marcus; Bood, Joakim

Published in:

Northern Optics & Photonics 2018, 12-14 September 2018, Lund University, Lund, Sweden

2018

Document Version:

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Ruchkina, M., Ding, P., Aldén, M., & Bood, J. (2018). Photolysis-free two photon laser induced fluorescence of H atom. In *Northern Optics & Photonics 2018, 12-14 September 2018, Lund University, Lund, Sweden: Conference Proceedings*

Total number of authors:

4

Creative Commons License:

Unspecified

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Photolysis-free two photon laser induced fluorescence of H atom

Maria Ruchkina^{1, *}, Pengji Ding¹, Marcus Aldén¹ and Joakim Bood¹

¹Div. of Combustion Physics, Department of Physics, Lund University, Box 118, SE-221 00 Lund, Sweden
email: maria.ruchkina@forbrf.lth.se
Telephone: 0722696428

Keywords: LIF; TPLIF; Two-photon excitation process; Femtosecond laser pulse; H atom

Abstract

Investigation of H atoms is highly demanded in the combustion environment since they are present in every chemical reaction which involves hydrocarbons. Optical detection of H atoms is difficult because the wavelength which is needed for the excitation is in the vacuum-UV region, where air is absorbing strongly. To overcome this problem two-photon excitation methods have been developed. Following the 2-photon excitation from $n=1$ to $n=3$ state the H atom emits fluorescence at the wavelength 656 nm. The use of fs-laser pulses benefits when it comes to the detection of species which might suffer from interfering photodissociation. The experiment was conducted where no difference in TPLIF (two-photon laser induced fluorescence) line profiles received for different laser output energies was found. This clearly indicates the absence of photolytically produced hydrogen atoms. TPLIF has many advantages, this technique allows imaging of atomic species in the premixed flame environment with high temporal and spatial resolution. The present investigation was conducted in a welding torch flame of 1.5 mm diameter, precisely at the tip of the flame. The experiment revealed that the concentration of H atoms at the top of the flame is lower than at the edges of the flame. It should be mentioned that LIF signal might suffer from SE (stimulated emission). Further investigations should be conducted in order to investigate the impact of SE on the signal.




Figure 1. TPLIF image of H atom fluorescence signal on a tip of CH₄/air flame ($\Phi = 1.3$), where the laser sheet is 7 mm height.