



LUND UNIVERSITY

Changes in Hygroscopicity and Cloud-Activation of Diesel Soot upon Ageing

Wittbom, Cerina; Svenningsson, Birgitta; Rissler, Jenny; Eriksson, Axel; Swietlicki, Erik; Nordin, Erik; Nilsson, Patrik; Pagels, Joakim

2012

[Link to publication](#)

Citation for published version (APA):

Wittbom, C., Svenningsson, B., Rissler, J., Eriksson, A., Swietlicki, E., Nordin, E., Nilsson, P., & Pagels, J. (2012). *Changes in Hygroscopicity and Cloud-Activation of Diesel Soot upon Ageing*. Abstract from Nordic Society for Aerosol Research (NOSA) Aerosol Symposium, 2012, Helsingör, Denmark.

Total number of authors:

8

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Changes in Hygroscopicity and Cloud-Activation of Diesel Soot upon Ageing

C. Wittbom¹, B. Svenningsson¹, J. Rissler², A. Eriksson¹, E. Swietlicki¹, E. Z. Nordin², P. T. Nilsson² and J. Pagels²

¹Department of Physics, Lund University, P.O. Box 118 SE 221 00, Lund, Sweden

²Ergonomics and Aerosol Technology, Lund University, P.O. Box 118 SE-221 00 Lund, Sweden

Keywords: H-TDMA, CCN, AMS, Organic compounds.

Introduction

The contribution from fresh diesel exhaust particles to the cloud condensation nuclei (CCN) population is negligible (e.g. Tritscher *et al.*, 2011). However, complex gas-to-particle conversion processes in the atmosphere form secondary organic aerosol (SOA) from emitted exhaust gases and particles, which significantly may influence the cloud drop formation process. Results from two Lund University smog chamber campaigns (#I & #II), are presented. The CCN and hygroscopic properties of diesel soot particles and the accompanying organic coating were investigated during ageing.

Methods

Exhaust from a light-duty diesel vehicle at warm idling were transferred to a smog chamber and photochemically aged (Nordin *et al.*, 2012). VOCs and IVOCs (Intermediate Volatile Organic Compounds) in the diesel exhaust were used as SOA precursors. Selected amounts of toluene and m-xylene were added to allow investigations of the full particle transformation from agglomerates to spheres.

Hygroscopic properties were analysed using a Hygroscopic Tandem Differential Mobility Analyzer (H-TDMA; Nilsson *et al.*, 2009), and the cloud-activation properties were measured using a Cloud Condensation Nucleus Counter (CCNC; DMT 100). A soot particle aerosol mass spectrometer (SP-AMS, Aerodyne Research) determined the composition of the soot cores and the particle coatings. The particle mass-mobility relationship was characterized using a Differential Mobility Analyzer-Aerosol Particle Mass Analyzer (DMA-APM; Kanomax Japan 3600). During Campaign II, the CCNC measurement procedure was changed from the traditional Stepping- ΔT to Scanning Flow CCN Analysis (SFCA) (Moore & Nenes, 2009), enabling rapid measurements of the supersaturation spectra with high time resolution, revealing more detailed, accurate and continuous results.

Conclusions

During the ageing process, the transformation of the hygroscopic behaviour and its link to the effect on cloud droplet activation were related to the organic fraction in the particle as well as particle size and morphology. The properties of the organic material strongly influence the activation of the coated soot particles, both regarding the change of chemical composition, and also the change of critical supersaturation (SS_c , Fig.1) upon UV exposure.




Figure 1. Ageing results in an increase in the mass growth factor (Gfm, x-axis), which is the ratio of total particle mass divided by the soot core mass. During ageing, the critical supersaturation (SS_c) decreases (red, left y-axis) and the corresponding hygroscopic growth factor (HGF, at 90% RH) increases (blue, right y-axis). Results are for diesel soot particles ($d_{m,dry}=150$ nm) from two experiments, Campaign #1 and #2.

Fresh diesel soot particles show no hygroscopic growth (HGF, Fig.1). Due to the morphology, the mobility diameter is not a relevant size measure for predicting cloud activation of fresh soot particles and is thereby not an accurate input parameter when calculating neither HGF (Fig.1) nor when modelling the SS_c . The cloud droplet activation starts long before the particles show any hygroscopic growth at all at 90% relative humidity (RH) (Fig.1). The soot particles gradually collapse to spheres up to a coating of 50-60% condensed organic material (HGF \approx 0.93; Gfm \approx 2-2.5, Fig.1). These results indicate that diesel soot particles, when aged, do affect the cloud forming process in the atmosphere.

The Swedish Research Council (VR), FORMAS and CRAICC supported this work.

Moore, R. & Nenes, A. (2009) *Aerosol Sci. & Tech.* 43:1192-1207.

Nilsson, E. et al. (2009) *Atmos. Meas. Tech.*, 2, 313-318.

Nordin, E.Z., et al. (2012) Smog chamber studies of SOA formation from Gasoline Vehicles & Aromatic precursors. *To be submitted to ACP*.

Tritscher, T., et al. (2011) *Environ. Res. Lett.* 6:3:034026.