



LUND UNIVERSITY

X-Ray Absorption in PIXE Analysis of Nonuniform Samples

Johansson, Gerd; Malmqvist, Klas; Carlsson, Lars-Eric; Akselsson, Roland

Published in:

Contribution to the Conference on Particle Induced X-Ray Emission and its Analytical Applications

1976

[Link to publication](#)

Citation for published version (APA):

Johansson, G., Malmqvist, K., Carlsson, L.-E., & Akselsson, R. (1976). X-Ray Absorption in PIXE Analysis of Nonuniform Samples. *Contribution to the Conference on Particle Induced X-Ray Emission and its Analytical Applications*.

Total number of authors:

4

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

X-Ray Absorption in PIXE Analysis of Nonuniform Samples

G. Johansson¹⁾, K. Malmqvist¹⁾, L.-E. Carlsson¹⁾ and
K.R. Akselsson^{1,2)}

- 1) Department of Nuclear Physics, Lund Institute of
Technology, Lund, Sweden
- 2) Department of Environmental Health, University of Lund,
Lund, Sweden

In PIXE analysis of small nonuniform samples corrections for X-ray absorption and slowing-down of protons may be very large and difficult to estimate. Inspection of a sample by microscope gives information about the shape but not about the density. However good estimations of the corrections for selfabsorption and proton slowing-down are obtained from measurements of the transmission of a narrow beam of low energy X-rays at several positions of the sample.

This paper demonstrates a feasible way of using this method. Welding aerosol samples from stage 5 of a Battelle designed cascade impactor are used as examples. An intense source of 3 keV X-rays is obtained by bombarding a thick silver target with 2 MeV protons. A 0.1 mm diameter X-ray beam is obtained by a pinhole collimator and the transmission of this beam in different parts of the sample is measured with a Si(Li)-detector.