



LUND UNIVERSITY

Extinction cross section measurements

Larsson, Christer; Sohl, Christian; Gustafsson, Mats; Kristensson, Gerhard

2008

[Link to publication](#)

Citation for published version (APA):

Larsson, C., Sohl, C., Gustafsson, M., & Kristensson, G. (2008). *Extinction cross section measurements*. Abstract from Svenska antenmätsällskapet, Arboga, Sweden.

Total number of authors:

4

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Extinction cross section measurements

Christer Larsson^{1,2}, Christian Sohl¹, Mats Gustafsson¹, and Gerhard Kristensson¹

¹Department of Electrical and Information Technology, Lund University,
P.O. Box 118, S-221 00 Lund, Sweden

²Saab Communication, S-581 11 Linköping, Sweden

Two methods to experimentally determine the extinction cross section for a large bandwidth in the microwave region are investigated in this paper. The motivation to measure the extinction cross section, *i.e.*, the sum of the total scattering cross section and the absorption cross section, comes from the need to verify recent theoretical results that bound the scattering from objects.

Through the optical theorem it is possible to calculate the extinction cross section from a measurement of the forward radar cross section (RCS). However, the direct measurement of the forward RCS in free space is experimentally difficult since the largest part of the received field at the receiving antenna consists of direct illumination by the transmitting antenna. The direct illumination contributes with a dominating background that has to be removed, with signal processing or otherwise, from the scattered field component that one wants to determine.

We have developed and validated a method to determine the extinction cross section for thin and non-magnetic planar objects. The method is based on a regular measurement of the monostatic RCS. The result shows that monostatic RCS measurements can be used with good accuracy to determine the extinction cross section for this type of thin samples.

The first method is compared to a more general measurement method based on a measurement of the RCS in the forward direction.

Examples will be given from measurements on fabricated samples of single-layer planar arrays of split ring resonators (SRR) or capacitive resonators. These designed materials are commonly described as metamaterials in the literature.