



LUND UNIVERSITY

Growth and respiration of aquatic bacteria on organic carbon from different terrestrial sources

Berggren, Martin; Laudon, Hjalmar; Jansson, Mats

2006

[Link to publication](#)

Citation for published version (APA):

Berggren, M., Laudon, H., & Jansson, M. (2006). *Growth and respiration of aquatic bacteria on organic carbon from different terrestrial sources*. Abstract from ASLO 2006 Summer Meeting - Global Challenges Facing Oceanography and Limnology, Victoria, B.C., Canada.

Total number of authors:

3

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Berggren M (oral), H Laudon and M Jansson. 2006. Growth and respiration of aquatic bacteria on organic carbon from different terrestrial sources. 2006 ASLO Summer Meeting: Global Challenges Facing Oceanography and Limnology, presented on June 9, 2006, Victoria, British Columbia, Canada

ABSTRACT

Streams and lakes are recipients for terrestrial drainage of organic carbon. This allochthonous organic carbon is metabolized, mainly by bacteria, with significant consequences for the biostructure and energy pathways of freshwater ecosystems. The degree to which these substrates support heterotrophic bacteria and their role as energy mobilizers is largely determined by bacterial growth efficiency (BGE), i.e., bacterial production per unit of assimilated carbon. We tested the hypothesis that catchment characteristics influence BGE by regulating the organic carbon quality and the availability of inorganic nutrients. Ten boreal headwater streams were examined (68°N 18°E), representing a gradient ranging from organic carbon supplied mainly from mires to carbon supplied mainly from forests. The forest coverage (%) of the catchment showed a positive correlation with BGE and BGE was positively correlated with the absorbance ratio A₂₅₄/365 and negatively with C:P and C:N ratios. The data suggest that there was a pool of rapidly utilized carbon which was drained mainly from forest soils and which was incorporated into bacterial biomass with great efficiency. Its exploitation was probably nutrient limited as indicated by nutrient enrichment experiments.