



LUND UNIVERSITY

Health Effects in Healthy Volunteers Exposed to Diesel Exhaust and Traffic Noise

Gudmundsson, Anders; Xu, Yiyi; Wierzbicka, Aneta; Andersson, Ulla B; Assarsson, Eva; Axmon, Anna; Barregård, Lars; Berglund, Margareta; Bohgard, Mats; Broberg Palmgren, Karin; Brunskog, Jonas; Gunnskog, Anna-Therese; Hagerman, Inger; Jönsson, Bo A; Nielsen, Jörn; Kåredal, Monica; Nilsson, Patrik; Österberg, Kai; Pagels, Joakim; Poulsen, Torben; Rissler, Jenny; Stockfelt, Leo; Sällsten, Gerd; Albin, Maria

Published in:

Proceedings of the "Workplace and Indoor Aerosols Conference" (electronic version only)

2012

[Link to publication](#)

Citation for published version (APA):

Gudmundsson, A., Xu, Y., Wierzbicka, A., Andersson, U. B., Assarsson, E., Axmon, A., Barregård, L., Berglund, M., Bohgard, M., Broberg Palmgren, K., Brunskog, J., Gunnskog, A.-T., Hagerman, I., Jönsson, B. A., Nielsen, J., Kåredal, M., Nilsson, P., Österberg, K., Pagels, J., ... Albin, M. (2012). Health Effects in Healthy Volunteers Exposed to Diesel Exhaust and Traffic Noise. In *Proceedings of the "Workplace and Indoor Aerosols Conference" (electronic version only)* (pp. 26-26)

Total number of authors:

24

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Health effects in healthy volunteers exposed to diesel exhaust and traffic noise.

Anders Gudmundsson¹, Yiyi Xu², Aneta Wierzbicka¹, Ulla B.K. Andersson², Eva Assarsson², Anna Axmon², Lars Barregård⁵, Margareta Berglund³, Mats Bohgard¹, Karin Broberg², Jonas Brunskog⁴, Anna-Therese Gunnskog², Inger Hagerman³, Bo A.G. Jönsson², Jörn Nielsen², Monica Kåredal², Patrik Nilsson¹, Kai Österberg², Joakim Pagels¹, Torben Poulsen⁴, Jenny Rissler¹, Leo Stockfelt⁵, Gerd Sällsten⁵, and Maria Albin²

¹Ergonomics and Aerosol Technology, Lund University, Sweden, ²Occupational and Environmental Medicine, Lund University, Sweden, ³Department of Cardiology, Karolinska University Hospital, Huddinge, Sweden,

⁴Acoustic Technology and Hearing Systems, Technical University of Denmark, Denmark, ⁵Occupational and Environmental Medicine, University of Gothenburg, Sweden

Presenting author email: anders.gudmundsson@design.lth.se

Introduction

Exposure to diesel exhaust (DE) and traffic noise is common. Previous studies found an inflammatory response in healthy volunteers, but no effect on lung function after one hour diesel exposure (particle mass concentration 350 µg/m³ and NO₂ concentration of 0.62 ppm).

Methods

Eighteen healthy volunteers were exposed in a stainless steel chamber four times, each 3 hours: 1) zero exposure, 2) high DE (300 µg/m³ and NO₂ 1.4 ppm) and low traffic noise, 3) low DE and high traffic noise 75 dB(A) and 4) high DE and high traffic noise.

Immediately before and after exposure we performed medical examination, spirometry, rhinometry and blood sampling (repeated next morning). Symptom scores and peak expiratory flow (PEF) and ECG measurements were assessed before, 15 min, 75 min and 135 min into exposure and after. Generalized Estimating Equation (GEE) model was used to analyze collected data.

Results

Self-rated irritation of the eyes and throat was higher during DE than non-diesel exposure (non-DE), with a statistically significant difference after 75 (eyes) and 135 min (eyes, throat). Signs of irritation in the upper airways were significantly more common after DE (OR=3.2) and tended to be so also for the eyes (OR=3.1, p=0.06). PEF increased during non-diesel exposure, but decreased during DE, with a statistically significant difference after 75

(+4.08 vs -9.58 l/s), and 135 min (+8.11 l/s vs -3.5 l/s).

Discussion

We found adverse acute effects with regard to symptoms, signs, PEF, and inflammatory markers in healthy volunteers exposed 3 hours to diesel (300 µg/m³ and NO₂ 1.4 ppm). TLV(8hours) of NO₂ is 1 ppm. These effects were first seen after 75 min of exposure. Preliminary results show correlations between heart rate variability and sound pressure.



WORKPLACE AND INDOOR AEROSOLS CONFERENCE

Lund University, 19-20 April 2012

PROCEEDINGS



The Swedish Research Council Formas
Committed to excellence in research for sustainable development



TRUST. SCIENCE. INNOVAT

